



The American Mathematical Monthly

ISSN: 0002-9890 (Print) 1930-0972 (Online) Journal homepage: www.tandfonline.com/journals/uamm20

# An Extension of Mantel's Theorem to k-Graphs

### Zoltán Füredi & András Gyárfás

To cite this article: Zoltán Füredi & András Gyárfás (2020) An Extension of Mantel's Theorem to *k*-Graphs, The American Mathematical Monthly, 127:3, 263-268, DOI: 10.1080/00029890.2020.1693227

To link to this article: https://doi.org/10.1080/00029890.2020.1693227

Published online: 24 Feb 2020.



Submit your article to this journal 🕝

Article views: 540



View related articles



View Crossmark data 🗹



Citing articles: 5 View citing articles 🕑

## An Extension of Mantel's Theorem to *k*-Graphs

#### Zoltán Füredi and András Gyárfás

**Abstract.** According to Mantel's theorem, a triangle-free graph on *n* points has at most  $n^2/4$  edges. A *linear k-graph* is a set of points together with some *k*-element subsets, called edges, such that any two edges intersect in at most one point. The *k*-graph  $F^k$ , called a *fan*, consists of *k* edges that pairwise intersect in exactly one point *v*, plus one more edge intersecting each of these edges in a point different from *v*. We extend Mantel's theorem as follows: fan-free linear *k*-graphs on *n* points have at most  $n^2/k^2$  edges.

This extension nicely illustrates the difficulties of hypergraph Turán problems. The determination of the case of equality leads to transversal designs on *n* points with *k* groups—for k = 3 these are equivalent to Latin squares. However, in contrast to the graph case, new structures and open problems emerge when *n* is not divisible by *k*.

**1. TRIANGLES AND FANS.** Once upon a time, 111 years ago, Mantel proposed a problem in a Dutch journal. He (followed by four other solvers) provided a solution and the outcome is known in graph theory as Mantel's theorem. Let  $K_p$  denote the complete graph on  $p \ge 2$  points;  $K_3$  is often called a triangle. Graphs without a  $K_p$  subgraph are called  $K_p$ -free.

#### **Theorem 1 (Mantel [4]).** Triangle-free graphs on n points have at most $n^2/4$ edges.

It is not easy to select the winner of the beauty contest for the nicest proof of Mantel's theorem. A visit to the internet brings proofs, some presented as videos [9]. A natural proof by induction competes with the proof of Mantel, using Cauchy's inequality. The book of Aigner and Ziegler [1] exhibits seven different proofs, five of them for Turán's generalization, the flagship theorem of extremal graph theory.

### **Theorem 2 (Turán [6]).** $K_p$ -free graphs on n points have at most $(1 - \frac{1}{p-1})\frac{n^2}{2}$ edges.

Analogues of Mantel's theorem are also considered for *k*-graphs, where the edges are *k*-element subsets of the points. Turán [7] asked about the maximum number of edges among 3-graphs on *n* points that contain no tetrahedron, four triples on four points. This is known as a notoriously difficult question, where even the asymptotic answer is unknown (conjectured to be  $\frac{5}{5} \binom{n}{3}$ ).

Here we have two aims. Namely, we present a more friendly extension of Mantel's theorem to k-graphs (Theorem 3). By doing so we illustrate many difficulties one can have encountering a Turán-type problem. Since many of these questions are unsolved, we investigate other important combinatorial structures between graphs and hypergraphs, like multigraphs (see, e.g., [3]), and linear hypergraphs. But even then, the solutions frequently lead to further unsolved problems.

A k-graph is called *linear* if any two edges intersect in at most one point. Note that graphs are linear 2-graphs. For any integer  $k \ge 2$ , the *fan*, denoted by  $F^k$ , is the k-graph

doi.org/10.1080/00029890.2020.1693227

MSC: Primary 05D05, Secondary 05B15; 05C70

having k + 1 edges,  $f_1, \ldots, f_k$ , and g, such that  $f_1, \ldots, f_k$  are disjoint apart from one common point v (*the center*) and an additional *crossing edge g* that intersects all  $f_i$  in points different from v. We prove the following extension of Mantel's theorem.

**Theorem 3.** Fan-free linear k-graphs on n points have at most  $n^2/k^2$  edges.

Considering all k-graphs, a different extension of Mantel's theorem arises.

**Theorem 4 (Mubayi and Pikhurko [5]).** Fan-free k-graphs on n > n(k) points have at most  $(n/k)^k$  edges.

*Proof of Theorem 3.* We denote by V(G), E(G) the set of points and edges of a *k*-graph *G*. The number of edges containing a point  $v \in V(G)$  is the *degree* of *v*, denoted by d(v). If all points have the same degree *m*, the *k*-graph is called *m*-regular.

Assume that *G* is a fan-free linear *k*-graph with *n* points. For any point  $v \in V(G)$ , set  $N_v := (\bigcup_{e \ni v} e) \setminus \{v\}$  (the open neighborhood of *v*), and  $B_v := V(G) \setminus N_v$ . Let  $\Delta = \Delta(G)$  be the maximum degree of *G* and select  $v \in V(G)$  such that  $d(v) = \Delta$ . Then  $|N_v| = (k-1)\Delta$  and  $|B_v| = n - (k-1)\Delta$ . Since *G* is fan-free, every edge of *G* must intersect  $B_v$ . This and the inequality  $d(x) \leq \Delta$  imply

$$|E(G)| \le \sum_{x \in B_v} d(x) \le \Delta |B_v| = \Delta(n - (k - 1)\Delta).$$
(1)

On the other hand, obviously

$$|E(G)| \le \frac{n\Delta}{k}.$$
(2)

If  $\Delta \leq (n/k)$  we immediately get  $|E(G)| \leq n^2/k^2$  from (2). If  $\Delta > (n/k)$  then (1) and the geometric–arithmetic mean inequality imply

$$|E(G)| \le \Delta(n - (k - 1)\Delta) \le \frac{1}{4}(n - (k - 2)\Delta)^2 < \frac{1}{4}\left(n - \frac{(k - 2)n}{k}\right)^2 = \frac{n^2}{k^2}.$$

Thus in both cases  $|E(G)| \le n^2/k^2$  as claimed.

**2. EXTREMAL CONFIGURATIONS.** It is interesting to see when we have equality in the bounds of the theorems of the previous section. Graphs (or *k*-graphs) attaining equality are called *extremal configurations*. In Theorem 2, the only extremal configuration is the balanced complete (p - 1)-partite graph: the points of *G* are divided into p - 1 groups, each with  $\frac{n}{p-1}$  points and we have all edges joining two points from different groups. In particular, for p = 3, the only extremal configuration in Theorem 1 is the balanced complete bipartite graph with even number of vertices. The extremal *k*-graphs of Theorem 3 must satisfy k|n; this is assumed in the next subsection.

**Transversal designs.** The complexity of extremal configurations grows with k; they are the transversal designs, defined as follows. Assume that n is a multiple of k. A *transversal design* T(n, k) is a linear k-graph on n points where the points are partitioned into k groups, each containing n/k points and where each pair of points from different groups belongs to exactly one edge. Note that for k = 2, T(n, 2) is the complete bipartite graph with n/2 points in its partite classes. The next case, k = 3, is

already more interesting. Assume that X, Y, Z are the groups of a transversal design T(n, 3) with points  $x_i \in X$ ,  $y_i \in Y$ ,  $z_i \in Z$  for i = 1, 2, ..., n/3. Consider the  $\frac{n}{3} \times \frac{n}{3}$  matrix A defined with  $a_{ij} = z_k$  where  $z_k$  is the (unique) point of Z for which  $x_i y_j z_k$  is an edge of T(n, 3). Each row and column of A contains different  $z_i$ 's; such a matrix is called a *Latin square* (of order n/3). For the interested reader we note that in general, transversal designs T(n, k) are equivalent to k - 2 mutually orthogonal Latin squares. The investigation of these combinatorial structures goes back to Euler. For a nice (and high-level) introduction, see van Lint and Wilson [8].

**Proposition 1.** Any T(n, k) is a fan-free linear k-graph with  $n^2/k^2$  edges.

*Proof.* Suppose that some T(n, k) contains a k-fan F with center v in group i. Since the crossing edge g of F must have a point in group i different from v, g can intersect at most k - 1 edges from the k edges of F containing v. This is a contradiction.

**Theorem 5.** Equality holds in Theorem 3 only if G is a transversal design T(n, k).

*Proof.* We continue the proof of Theorem 3. If equality holds, then inequalities (1) and (2) are equalities. From (2) we have that n = km and G is an *m*-regular *k*-graph. It is left to show that  $|E(G)| = m^2$  implies that G is a transversal design with k groups. Since G is *m*-regular, we have  $|B_v| = km - (k - 1)m = m$  for every  $v \in V(G)$ .

**Claim 1.** For every  $v \in V(G)$ ,  $B_v$  is a strongly independent set, i.e., every edge intersects it in at most one point.

To prove the claim, assume that  $x, y \in B_v$  and that there is an edge  $e \in E(G)$  containing x, y. Then (1) cannot be an equality, since e is counted from both x and y. This is a contradiction, proving the claim.

Applying the claim for the points of an arbitrary edge  $e = \{v_1, v_2, \ldots, v_k\}$ , we get the strongly independent sets  $B_{v_1}, \ldots, B_{v_k}$ . These sets must be pairwise disjoint, because if  $x \in B_{v_i} \cap B_{v_j}$  then  $B_{v_j} \cup \{v_i\} \subseteq B_x$ , contradicting the fact that  $|B_{v_j}| = |B_x|$ . Thus V(G) can be partitioned into k groups of size m, each forming a strongly independent set. The  $m^2$  edges of G cover  $m^2 {k \choose 2}$  pairs in V(G) and this is equal to  ${mk \choose 2} - k{m \choose 2}$ , the number of pairs of V(G) not covered by the groups  $B_{v_1}, \ldots, B_{v_k}$ . Thus each pair of points from different groups is covered exactly once, proving that G is a transversal design with k groups of size m.

**Truncated designs.** What happens when *n* is not divisible by *k*? Turán [6] proved that the unique extremal configuration for  $K_p$ -free graphs on *n* points is the following graph (the Turán-graph): *n* points are divided into p - 1 groups as evenly as possible and the edges are all pairs of points from different groups. Considering the same question for fan-free *k*-graphs, we can answer only in the case  $n \equiv -1 \pmod{k}$  for general *k*, as far as the maximum number of edges is concerned. A *truncated design* is obtained from a transversal design by removing one point (and all edges containing it).

**Theorem 6.** Assume that  $k \ge 2$ , n = k(m + 1) - 1, and G is a fan-free k-graph with n points. Then  $|E(G)| \le m^2 + m$ . Truncated designs obtained from T(n + 1, k) are extremal configurations.

*Proof.* It follows from Proposition 1 that any truncated design obtained from T(n + 1, k) is a fan-free linear k-graph with  $m^2 + m$  edges. To show that  $|E(G)| \le m^2 + m$ 

March 2020]

whenever G is a fan-free linear k-graph with n = km + k - 1 points, we follow the argument of the proof of Theorem 3 using the same notations.

If  $\Delta \leq m$ , we immediately get from (2) that

$$|E(G)| \le \frac{((m+1)k-1)m}{k} = m^2 + m - \frac{m}{k} < m^2 + m.$$

If  $\Delta \ge m + 1$ , then (1) and the geometric–arithmetic mean inequality imply

$$|E(G)| \le \Delta(n - (k - 1)\Delta) \le \frac{1}{4}(n - (k - 2)\Delta)^2$$
$$\le \frac{1}{4}((m + 1)k - 1 - (k - 2)(m + 1))^2 = m^2 + m + \frac{1}{4}$$

Thus in both cases  $|E(G)| \le m^2 + m$ .

**3. EXTREMAL TRIPLE SYSTEMS.** In this section, we refer to linear 3-graphs as triple systems.

**Extensions of triangle-free graphs.** To find all extremal configurations for k = 3,  $n \equiv -1 \pmod{3}$ , we need a special case of a theorem of Andrásfai, Erdős, and Sós, stated here as a lemma.

**Lemma 1** (Andrásfai, Erdős, and Sós [2]). Assume that a nonbipartite graph G has n points and contains no triangles. Then the minimum degree of G is at most 2n/5.

The following graph, *the blown up five-cycle*,  $C_5^t$ , shows that Lemma 1 is sharp when  $n \equiv 0 \pmod{5}$ . Take a five-cycle and replace its points with disjoint *t*-element sets of points,  $A_1, \ldots, A_5$  and replace its edges by complete bipartite graphs  $[A_i, A_{i+1}]$  for  $i = 1, \ldots, 5 \pmod{5}$ .

There is an easy way to generate fan-free triple systems from triangle-free graphs. Consider a graph G with a proper edge-coloring, i.e., let the edge set of G be partitioned into d matchings (pairwise disjoint edges)  $M_1, \ldots, M_d$ . The *extension* of G, T(G), is the triple system obtained by extending V(G) with d new points  $v_1, \ldots, v_d$  and extending every edge of  $M_i$  with the point  $v_i$ , for  $i = 1, \ldots, d$ .

**Proposition 2.** Assume that G is a triangle-free graph with a proper edge coloring. Then T(G) is a fan-free triple system.

*Proof.* No fan in T(G) can be centered at  $v_i$  since three edges of  $M_i$  cannot be intersected by any edge of  $M_j$  for  $j \neq i$ . If a fan in T(G) is centered at  $w \in V(G)$  with triples  $ww_1v_1, ww_2v_2, ww_3v_3$ , then its crossing edge must be of the form  $w_iw_jv_k$  with three different indices. Thus  $w, w_i, w_j$  is a triangle in G, contradicting the assumption.

An important special case of proper edge colorings is the 1-factorization, where all of the d matchings cover all points of the graph. Graphs having 1-factorizations are obviously d-regular but the converse statement is not true: odd cycles are easy examples. The most famous example is the *Petersen graph*.

We will apply Proposition 2 to two triangle-free nonbipartite graphs. One of them is the Wagner graph,  $C_8^*$ , the eight-cycle with its long diagonals; the other is  $C_5^2$ , defined earlier in this section.

#### Extremal triple systems for $n \equiv -1 \pmod{3}$ .

**Theorem 7.** Assume that G is a linear fan-free triple system with n = 3m + 2 points. Then G has at most  $m^2 + m$  edges. Equality holds only in the following cases.

- (7.1) *G* is a truncated design obtained from a transversal design T(3m + 3, 3),
- (7.2) m = 3, G is the extension of a 1-factorization of  $C_8^*$ ,
- (7.3) m = 4, G is the extension of a 1-factorization of  $C_5^2$ .

*Proof.* We use the notation of the proof of Theorem 6. Assume that G is a fan-free linear 3-graph, |V(G)| = 3m + 2,  $|E(G)| = m^2 + m$ .

From (2) it follows that  $\Delta \ge m + 1$ . The inequality (1) would give  $|E(G)| < m^2 + m$  if any vertex of  $v \in V(G)$  has degree larger than m + 1. Thus  $\Delta(H) = m + 1$ . Suppose  $d(x) = m + 1 = \Delta$ . Then for  $B := B_x$  one has |B| = m, all points of  $B_x$  have degree m + 1, and  $B_x$  is a strongly independent set. It follows that  $N_y = N_x$  for each  $y \in B$ , implying  $d_H(w) = m$  for all  $w \in W := V(G) \setminus B$ .

Define the graph  $G^*$  on point set W, where  $w_1, w_2 \in W$  is an edge in  $G^*$  if and only if  $w_1w_2v$  is an edge of the 3-graph G for some  $v \in B$ . Then  $G^*$  is *m*-regular and can be written as the union of *m* matchings of size m + 1. If  $G^*$  is bipartite, then it must be isomorphic to  $K_{m+1,m+1}$  with one matching removed; thus G is a truncated design obtained from three groups of size m + 1, the first possibility in Theorem 7.

We claim that  $G^*$  is a triangle-free graph. Indeed, if  $w_1, w_2, w_3$  form a triangle in  $G^*$ , then G contains the edges

$$e = w_1 w_2 v_3, f = w_1 w_3 v_2, g = w_2 w_3 v_1$$

for  $v_i \in B_v$ . Because  $d_G(v_1) = m + 1$ , there is an edge  $h = v_1 w_1 w_4$ . Then e, f, g, h form a fan with center  $w_1$  and with crossing edge g, a contradiction.

Thus we may suppose that  $G^*$  is a nonbipartite triangle-free graph. Applying Lemma 1 to our graph  $G^*$ , we get  $m \le 2(2m+2)/5$ ; thus  $m \le 4$ . We leave it to the reader to check that there are no *m*-regular nonbipartite triangle-free graphs on 2m + 2 points for m = 1, 2, but for m = 3, 4 there are unique ones:  $C_8^*, C_5^2$ . Moreover, both graphs have 1-factorizations. Extending them with the points of *B*, we get the second and third possibilities in Theorem 7.

**Remark.** Theorem 7 seemingly provides two exceptional extremal configurations. However, this is not right: there are *three*! The explanation is that the Wagner graph has two nonisomorphic 1-factorizations (and  $C_5^2$  has only one).

**4. CONCLUSION.** It does not seem easy to find, for every  $n \neq 0 \pmod{k}$ , the maximum number of edges in fan-free linear *k*-graphs on *n* points, let alone give a description of all extremal *k*-graphs attaining this maximum. We could answer the former question only in the case  $n \equiv -1 \pmod{k}$  in Theorem 6 and the latter in the subcase k = 3 in Theorem 7. The study of the remaining cases might reveal some (possibly infinitely many) exceptional extremal configurations. We *conjecture* that the extremal number is  $m^2$  for m > m(k) and  $n \neq -1 \pmod{k}$ .

We conclude with a *conjecture* that generalizes Theorems 3 and 4 as well: If n = km, m > m(k),  $1 \le \ell \le k$ , then an  $F^k$ -free k-graph on n points with more than  $m^\ell$  edges has two edges intersecting in at least  $\ell$  points.

ACKNOWLEDGMENT. The authors are grateful for the helpful comments of a referee.

March 2020]

- [1] Aigner, M., Ziegler, G. (2004). Proofs from THE BOOK. Berlin-Heidelberg: Springer-Verlag.
- [2] Andrásfai, B., Erdős, P., Sós, V. T. (1974). On the connection between chromatic number, maximal clique and minimal degree of a graph. *Discrete Math.* 8(3): 205–218.
- [3] Füredi, Z., Kündgen, A. (2002). Turán problems for integer-weighted graphs. J. Graph Theory. 40(4): 195–225.
- [4] Mantel, M. (1907). Problem 28. Wiskundige Opgaven. 10: 60-61.
- [5] Mubayi, D., Pikhurko, O. (2007). A new generalization of Mantel's theorem to k-graphs. J. Comb. Theory Ser. B. 97(4): 669–678.
- [6] Turán, P. (1941). On an extremal problem in graph theory (in Hungarian). Mat. Fiz. Lapok. 48: 436–452.
- [7] Turán, P. (1961). Research problem. Közl. MTA Mat. Kutató Int. 6: 177–181.
- [8] van Lint, J. H., Wilson, R. M. (2001). *A Course in Combinatorics*, 2nd ed. Cambridge: Cambridge Univ. Press.
- [9] Mantel's theorem. www.youtube.com/watch?v=WgddUjZVyj4

Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences furedi.zoltan@renyi.mta.hu

Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences gyarfas.andras@renyi.mta.hu