# Turán number of special four cycles in triple systems

Zoltán Füredi, András Gyárfás, Attila Sali

Alfréd Rényi Institute of Mathematics Hungarian Academy of Sciences

November 19, 2021

#### Abstract

A special four-cycle F in a triple system consists of four triples inducing a  $C_4$ . This means that F has four special vertices  $v_1, v_2, v_3, v_4$  and four triples in the form  $w_i v_i v_{i+1}$  (indices are understood (mod 4)) where the  $w_j$ s are not necessarily distinct but disjoint from  $\{v_1, v_2, v_3, v_4\}$ . There are seven non-isomorphic special four-cycles, their family is denoted by  $\mathcal{F}$ . Our main result implies that the Turán number  $ex(n, \mathcal{F}) = \Theta(n^{3/2})$ . In fact, we prove more,  $ex(n, \{F_1, F_2, F_3\}) = \Theta(n^{3/2})$ , where the  $F_i$ -s are specific members of  $\mathcal{F}$ . This extends previous bounds for the Turán number of triple systems containing no Berge four cycles.

We also study  $ex(n, \mathcal{A})$  for all  $\mathcal{A} \subseteq \mathcal{F}$ . For 16 choices of  $\mathcal{A}$  we show that  $ex(n, \mathcal{A}) = \Theta(n^{3/2})$ , for 92 choices of  $\mathcal{A}$  we find that  $ex(n, \mathcal{A}) = \Theta(n^2)$  and the other 18 cases remain unsolved.

## 1 Introduction

A triple system H = (V, E) has vertex set V and E consists of some triples of V (repeated triples are excluded). For any fixed family  $\mathcal{H}$  of triple systems, the Turán number  $ex(n, \mathcal{H})$  is the maximum number of triples in a triple system of n vertices that is  $\mathcal{H}$ -free, i.e., does not contain any member of  $\mathcal{H}$  as a subsystem.

Our interest here is the family  $\mathcal{F}$  of special four cycles: they have four distinct base vertices  $v_1, v_2, v_3, v_4$  and four triples  $w_i v_i v_{i+1}$  (indices are understood (mod 4)) where the  $w_j$ s are not necessarily distinct but  $w_i \neq v_j$  for any pair of indices  $1 \leq i, j \leq 4$ .

There are seven non-isomorphic special four cycles. The linear (loose) four cycle  $F_1$  is obtained when all  $w_j$ -s are different and in  $F_2$  all  $w_j$ s coincide. When two pairs coincide we get either  $F_3$  ( $w_1 = w_2, w_3 = w_4$ ) or  $F_4$  ( $w_1 = w_3, w_2 = w_4$ ). The  $F_4$  is the Pasch configuration. We define  $F_5$  with  $w_1 = w_2 = w_3$  (but  $w_4$  is different). In  $F_6$ 



Figure 1: The family of special four cycles  $F_1, F_2, F_3$  of Theorem 1



Figure 2: The family of the other four special four cycles  $F_4, \ldots, F_7$ 

we have  $w_1 = w_3$  (and  $w_2, w_4$  are different from  $w_1$  and from each other). When only  $w_1, w_2$  coincide we get  $F_7$ . Set  $\mathcal{F} = \{F_1, \ldots, F_7\}$ . For the convenience of the reader, the special four cycles are shown on Figure 1 and Figure 2.

Turán numbers of various members of  $\mathcal{F}$  have been investigated before. Füredi [4] proved that  $ex(n, F_3) \leq \frac{7}{2} \binom{n}{2}$ . Mubayi [7] showed that  $ex(n, F_2) = \Theta(n^{5/2})$ . Rödl and Phelps [8] gave the bounds  $c_1 n^{5/2} \leq ex(n, F_4) \leq c_2 n^{11/4}$ . In fact, the upper bound is Erdős' upper bound [2] for  $ex(n, K_{2,2,2}^3)$ . The lower bound comes from a balanced 3-partite triple system where every vertex of the third partite class form a triple with the edges of a bipartite  $C_4$ -free graph between the first two partite classes.

We prove that  $ex(n, \mathcal{F}) = \Theta(n^{3/2})$ , thus has the same order of magnitude as  $ex(n, C_4)$  for graphs. In fact, it is enough to exclude three of the special four cycles.

## **Theorem 1.** $ex(n, \{F_1, F_2, F_3\}) = \Theta(n^{3/2}).$

The family  $\mathcal{F}$  of special four cycles is a subfamily of a wider class, the class of *Berge four cycles*, where the vertices  $w_i$  can be selected from the base vertices as well, requiring only that the four triples  $w_i v_i v_{i+1}$  are different. Theorem 1 extends previous similar upper bounds (Füredi and Özkahya [5], Gerbner, Methuku, Vizer [6]) where the family of Berge four cycles were forbidden.

The appearance of the set  $\{F_1, F_2, F_3\}$  is not accidental. If any of  $F_1, F_2, F_3$  is missing from  $\mathcal{A} \subset \mathcal{F}$  then  $ex(n, \mathcal{A})$  is essentially larger than  $n^{3/2}$ .

- (C1) Ruzsa and Szemerédi [9] constructed triple systems on n vertices that do not carry three triples on six vertices and have more than  $n^{2-\varepsilon}$  triples for any fixed  $\varepsilon$ . This provides an example which contains only  $F_1$  from  $\mathcal{F}$ ,
- (C2) The  $\binom{n-1}{2}$  triples containing a fixed vertex from n vertices contains only  $F_2$  from  $\mathcal{F}$ ,
- (C3) Partition n vertices evenly into three parts, take a pairing between two equal parts and extend each pair with all vertices of the third class to a triple. This gives a triple system with approximately  $n^2/9$  triples and contains only  $F_3$  from  $\mathcal{F}$ .

In Section 3 we discuss  $ex(n, \mathcal{A})$  for all  $\mathcal{A} \subseteq \mathcal{F}$ . It turns out that in 92 cases  $ex(n, \mathcal{A}) = \Theta(n^2)$  and 18 cases remain unsolved.

## 2 Proof of Theorem 1

Assume *H* is a triple system with *n* vertices containing no subsystem from the set  $F_1, F_2, F_3$ . Applying the standard approach (based on [3]), we may assume that *H* is 3-partite with vertex partition  $[A_1, A_2, A_3]$  where  $|A_i| \in \{\lfloor n/3 \rfloor, \lceil n/3 \rceil\}$  and contains at least 2/9 of the triples of the original triple system.

The triples of H define a bipartite graph  $B = [A_1, A_2]$  as follows. If  $(a_1, a_2, a_3)$  is a triple of H with  $a_i \in A_i$  then  $a_1a_2$  is considered as an edge of B. Define the label  $L(a_1, a_2)$  of  $a_1a_2 \in E(B)$  as the set  $\{z \in A_3 : a_1a_2z \in E(H)\}$ . Then

$$|E(H)| = \sum_{a_1 a_2 \in E(B)} |L(a_1, a_2)|.$$
(1)

**Lemma 1.** The bipartite graph B has at most  $O(n^{3/2})$  edges.

**Proof of Lemma 1.** We denote by N(x, y) the set of common neighbors (in B) of  $x, y \in A_2$  in  $A_1$ . Similarly, let N(u, v) be the set common neighbors of  $u, v \in A_1$  in  $A_2$ .

For distinct vertices  $x, y \in A_2$ , define the digraph D = D(x, y) with vertex set  $A_3$ . For every  $u \in A_1$  such that  $u \in N(x, y)$  and  $a_i \in L(u, x), a_j \in L(u, y)$ , a directed edge  $a_i a_j$  is defined in D(x, y). We claim that D(x, y) is a very special digraph.

#### Claim 1.

- (1.1) There are no multiple loops or parallel directed edges in D(x, y),
- (1.2) There is at most one loop in D(x, y),

• (1.3) Two non-loop edges of D(x, y) either intersect or  $|N(x, y)| \le 4$ .

**Proof.** A multiple loop  $a_i a_i$  in D(x, y) would give a  $C_4$  in B with all edges containing  $a_i$  in their labels, this corresponds to an  $F_2$  in H – a contradiction. A multiple edge  $a_i a_j$  would give a  $C_4 = (x, u_1, y, u_2)$  in B where  $u_1, u_2 \in N(x, y), u_1 \neq u_2$  such that the consecutive edges of  $C_4$  contain  $a_i, a_j, a_j, a_i \in A_3$  in their labels. This would give an  $F_3$  in H – a contradiction again, proving (1.1).

Two distinct loops  $a_i a_i, a_j a_j$  in D(x, y) can appear in two ways. Either we have a  $C_4 = (x, u_1, y, u_2)$  in B where  $u_1, u_2 \in N(x, y), u_1 \neq u_2$  such that the consecutive edges of  $C_4$  contain  $a_i, a_i, a_j, a_j \in A_3$  in their labels, this would give an  $F_3$  in H, a contradiction. Otherwise  $u = u_1 = u_2$  and we have two multiedges xu, yu both containing  $a_i, a_j$  in their labels, this gives an  $F_2$  in H with u in its center, a contradiction again, proving (1.2).

Suppose that there exists two non-intersecting non-loop edges  $a_i a_j$ ,  $a_k a_l$  in D(x, y). If these edges are defined by  $u_1, u_2 \in N(x, y), u_1 \neq u_2$ , we have a  $C_4 = (x, u_1, y, u_2)$ in B with four distinct elements in their labels, giving an  $F_1$  in H, a contradiction. Thus we may assume that  $u_1 = u_2 = u$  and we have xu, yu in B with  $a_i, a_k$  and with  $a_j, a_l$  in their labels. Set

$$M = \{ v \in N(x, y) : v \neq u, |L(v, x) \cup L(v, y)| \ge 2 \}.$$

We claim that  $|M| \leq 2$ . Indeed, consider  $v \in M$ , there is  $a_s, a_t \in A_3$  such that  $a_s \neq a_t$  and xv, yv have labels containing  $a_s, a_t$ , respectively. Observe that either  $\{s,t\} = \{i,k\}$  or  $\{s,t\} = \{j,l\}$  otherwise there is a  $C_4 = (x, u, y, v)$  with four distinct labels, giving an  $F_1$  in H, a contradiction. This implies that  $|M| \leq 4$ . However, it cannot happen that for two distinct vertices  $v, v' \in M$  the coincidence of the index pairs are  $\{i,k\}$  and  $\{j,l\}$ , respectively, because it would result again in a  $C_4 = (x, v, y, v')$  with four distinct labels, a contradiction as above. Thus  $|M| \leq 2$  (equality is possible with edge pairs  $a_i a_k, a_k a_i$  or  $a_j a_l, a_l a_j$ ), proving the claim.

Observing that every vertex of  $N(x, y) \setminus (\{u\} \cup M)$  defines a loop in D(x, y), (1.1) and (1.2) implies that  $|N(x, y)| \leq 4$ , proving (1.3) and Claim 1.  $\Box$ 

A cherry on  $x \in A_2$  is defined as an incident edge pair,  $ux, vx \in E(B)$  such that  $u, v \in A_1, u \neq v$  and  $L(u, x) \cap L(v, x) \neq \emptyset$ . Let C(x, y) be the number of cherries in the subgraph of B induced on  $\{x, y\} \cup N(x, y)$ . We claim

$$C(x,y) \ge \sum_{a \in V(D(x,y))} d^+(a) + d^-(a) - 2,$$
(2)

because there are at least  $d^+(a) - 1$  cherries on x with  $L(u, x) \cap L(v, x) = \{a\}$  and  $d^-(a) - 1$  cherries on y with  $L(u, y) \cap L(v, y) = \{a\}$ .

Claim 2. For any two distinct vertices  $x, y \in A_2$ ,  $C(x, y) \ge |N(x, y)| - 4$ .

**Proof.** It is certainly true for  $|N(x, y)| \leq 4$ . Otherwise, using (1.3) from Claim 1, we have pairwise intersecting edges in D(x, y).

**Case 1.** The edges of D(x, y) form a triangle (edges oriented two ways are allowed) plus at most one loop. Therefore D(x, y) has at most seven edges thus  $5 \le |N(x, y)| \le$ 7. by (1.1) of Claim 1,  $d^+(a) \le 2$  for any vertex of the triangle. There are at least |N(x, y)| - 1 edges on the triangle, so there exists at least |N(x, y)| - 4 vertices a with  $d^+(a) \ge 2$  resulting in at least |N(x, y)| - 4 cherries on x.

**Case 2.** All edges of D(x, y) (apart from a possible loop) contain  $a \in A_3$ . For every  $u \in N(x, y)$  (apart from one possible vertex which defines a loop) either ux or uy has label a. Thus  $\sum_{a \in V(D(x,y))} d^+(a) + d^-(a) \ge |N(x,y)| - 1$  so (2) results in at least |N(x,y)| - 3 cherries on x or on y, completing the proof of Claim 2.  $\Box$ 

Claim 3. 
$$\sum_{x,y\in A_2} C(x,y) \leq {\binom{|A_1|}{2}}.$$

**Proof.** Every cherry counted on the left hand side is on some pair of  $A_1$ . At most one cherry can be on any  $(u, v) \in A_1$ , otherwise (by (1.1) in Claim 1) we have one of  $F_2, F_3$ .  $\Box$ 

Applying Claims 2, 3 we get

$$\sum_{x,y\in A_2} (|N(x,y)| - 4) \le \sum_{x,y\in A_2} C(x,y) \le \binom{|A_1|}{2},$$

thus

$$\sum_{x,y\in A_2} |N(x,y)| \le 4 \binom{|A_2|}{2} + \binom{|A_1|}{2} \le O(n^2).$$

By convexity we get

$$|A_1|\binom{|E(B)|}{|A_1|}{2} \le \sum_{u \in A_1} \binom{d(u)}{2} = \sum_{x,y \in A_2} |N(x,y)| \le O(n^2),$$

therefore  $|E(B)| = O(n^{3/2})$ , proving Lemma 1.  $\Box$ 

To finish the proof of Theorem 1, we need to show that the presence of labels does not affect strongly the edge count of Lemma 1. Let  $B^*$  denote the subgraph of Bwith the edges of at least three-element labels.

**Proposition 1.** If H is  $\{F_1, F_2, F_3\}$ -free then  $B^*$  is  $C_4$ -free.

**Proof.** Assume C = (x, u, y, v, x) is a four-cycle in  $B^*$ . From the definition of  $B^*$  there are three distinct elements, say a, b, c from the labels of three edges of C. The

only way to avoid  $F_1$  is that the fourth edge has label  $\{a, b, c\}$ . However, the same argument forces that all labels on C are equal to  $\{a, b, c\}$  giving (many)  $F_3$ 's.  $\Box$ 

We can consider  $B^*$  as a bipartite multigraph obtained as the union of  $|A_3|$  simple bipartite graphs as follows. Set

$$E(z) = \{(u, x) : u \in A_1, x \in A_2, (u, x, z) \in E(H) \text{ and } |L(u, x)| \ge 3\},\$$

then  $E(B^*) = \bigcup_{z \in A_3} E(z)$ .

**Proposition 2.** For every  $z \in A_3$  there is no path in  $B^*$  with four edges such that its first and last edge is in E(z).

**Proof.** Suppose that edges  $e_1, e_2, e_3, e_4$  form such a path for some  $z \in A_3$ . Since each edge of  $B^*$  has multiplicity at least three, we can replace  $e_2$  by  $f_2$  and  $e_3$  by  $f_3$  so that  $f_2 \in E(z_1), f_3 \in E(z_2)$  and  $z_1, z_2$  are distinct and both different from z. Then the four triples of H,

$$e_1 \cup \{z\}, f_2 \cup \{z_1\}, f_3 \cup \{z_2\}, e_4 \cup \{z\}$$

form an  $F_1$ , contradiction.  $\Box$ 

For any vertex  $x \in A_2$  let L(x) denote the subset of  $A_3$  that appears in some of the labels on edges of  $B^*$  incident to x.

**Proposition 3.** For distinct vertices  $x_1, x_2, x_3, x_4 \in A_2$ ,

$$|L(x_1) \cap L(x_2) \cap L(x_3) \cap L(x_4)| \le 1.$$

**Proof.** Suppose on the contrary that we have  $z_1, z_2 \in A_3$  such that for i = 1, 2, 3, 4,  $e_i = \{z_1, x_i, u_{2i-1}\}, f_i = \{z_2, x_i, u_{2i}\}$  are all triples of *H*.

An  $F_1$  is formed by the triples  $e_i, f_i, e_j, f_j$  if there is a pair i, j such that  $u_{2i-1}, u_{2i}, u_{2j-1}, u_{2j}$  are all different. Thus, we may assume that for any pair  $1 \le i < j \le 4$  there is an equality between elements  $u_{2i-1}, u_{2i}, u_{2j-1}, u_{2j}$ .

Let us call an equality  $u_{2i-1} = u_{2i}$  horizontal, an equality  $u_{2i} = u_{2j}$  or  $u_{2i-1} = u_{2j-1}$ (for  $i \neq j$ ) vertical, finally an equality  $u_{2i-1} = u_{2j}$  (for  $i \neq j$ ) diagonal. The terms to distinguish equalities refer to an arrangement of the vertices  $u_i$  into a  $4 \times 2$  matrix with  $u_{2i-1}, u_{2i}$  in row *i*. Observe the following facts.

- 1.  $F_3$  or  $F_2$  is formed by the triples  $e_i, f_i, e_j, f_j$  if the pair  $i \neq j$  have both horizontal equalities holding. Thus, at most one horizontal equality may hold.
- 2. If there is pair  $i \neq j$  such that both vertical equalities hold, then a  $C_4$  can be found in  $B^*$  contradicting to Proposition 1. Similarly,

- 3. if there is pair  $i \neq j$  such that both diagonal equalities hold, we get a contradiction with Proposition 1.
- 4. We get a four edge path contradicting to Proposition 2 if there are is a pair i, jsuch that exactly one vertical equality holds, that is  $u_{2i} = u_{2j}$  and  $u_{2i-1}, u_{2j-1}$ are different and different from  $u_{2i}$  as well. (Symmetrically, if there are  $x_i, x_j$ such that  $u_{2i-1} = u_{2j-1}$  and  $u_{2i}, u_{2j}$  are different and different from  $u_{2i-1}$  as well.)

Facts 1–4 imply that there exists a triple of indices, i, j, k such that we have exactly one diagonal equality on each pair of them. These are either in the form  $u_{2i-1} = u_{2k}, u_{2i} = u_{2j-1}, u_{2j} = u_{2k-1}$  defining a six-cycle in B on the vertices  $x_i, x_j, x_k, u_{2i}, u_{2j}, u_{2k}$ , giving (three)  $F_1$ , for example  $e_i, f_k, f_j, e_j$ , or in the form  $u_{2i-1} = u_{2k}, u_{2i} = u_{2j-1}, u_{2j-1} = u_{2k}$  that implies horizontal equality  $u_{2i-1} = u_{2i}$ , a contradiction. This proves Proposition 3.  $\Box$ 

By Propositions 1, 2 the simple bipartite graph B(z) with edge set E(z) has no cycles or paths with four edges. Therefore each component of B(z) is a double star. Thus each B(z) can be written as the union of two graphs, S(z), T(z) where each vertex of  $S(z) \cap A_2$  and each vertex of  $T(z) \cap A_1$  has degree one in B(z). Set

$$S = \bigcup_{z \in A_3} S(z), T = \bigcup_{z \in A_3} T(z).$$

By the definition of S, for every vertex  $x \in A_2$ , we have  $|L(x)| = d_S(x)$  where  $d_S(x)$  is the degree of vertex x in the (multi) graph S. By Proposition 3

$$\sum_{x \in A_2} \binom{|L(x)|}{2} \le 3\binom{|A_3|}{2}$$

therefore

$$\sum_{x \in A_2} \binom{d_S(x)}{2} \le 3 \binom{|A_3|}{2}.$$

Applying the same argument symmetrically for vertices of  $A_1$  and for the graph T, we get

$$\sum_{u \in A_1} \binom{d_T(u)}{2} \le 3 \binom{|A_3|}{2}.$$

By the convexity argument,  $|E(B^*)| = |E(S)| + |E(T)| = O(n^{3/2})$ . By Lemma 1, we also have  $|E(B)| = O(n^{3/2})$ . Thus by (1) and the definition of  $B^*$ ,

$$|E(H)| = \sum_{a_1 a_2 \in E(B)} |L(a_1, a_2)| \le 2|E(B)| + |E(B^*)| = O(n^{3/2}),$$

concluding the proof of Theorem 1.

## **3** Concluding remarks

Theorem 1 determines the order of magnitude  $(\Theta(n^{3/2}))$  for the 16 subsets of  $\mathcal{F}$  containing  $F_1, F_2, F_3$  and we pointed out that for all other choices  $\mathcal{A} \subset \mathcal{F}$ ,  $ex(n, \mathcal{A})$  must be essentially larger. In this section we summarize what we know about these cases. There is a trivial case, when  $\mathcal{A}$  is empty and  $ex(n, \mathcal{A}) = {n \choose 3}$ . Furthermore, as mentioned before,  $ex(n, F_2) = \Theta(n^{5/2})$  was proved by the first author (see in Mubayi [7]). Thus we have  $2^7 - 2^4 - 2 = 110$  cases to consider. It turns out that in 92 cases the order of magnitude is  $\Theta(n^2)$  (see Subsection 3.1) and only the remaining 18 cases are left unsolved (see Subsection 3.2).

A simple but useful lemma compares Turán numbers of closely related triple systems. Assume G is a triple system and  $v, w \in V(G)$  is covered by  $e \in E(G)$ . The triple system obtained from G by removing e and adding the triple v, w, x where  $x \notin V(G)$  is called a *fold out* of G. For example  $F_7$  is a fold out of  $F_3$ ,  $F_6$  is a fold out of  $F_4$ .

**Lemma 2.** (Fold out lemma.) If G is a triple system and  $G_1$  is a fold out of G then  $ex(n, G_1) \leq ex(n, G) + (|V(G)| - 2) {n \choose 2}$ .

**Proof.** Suppose that a triple system H has n vertices and has more than  $ex(n, G) + (|V(G)| - 2)\binom{n}{2}$  triples. A triple of H is called bad if it contains a pair of vertices that covered by at most |V(G)| - 2 triples of H, otherwise it is a good triple. Then H has more than ex(n, G) good triples thus contains a copy of G with all triples good. By definition, any pair of vertices in any triple of this copy of G is in more than |V(G)| - 2 triples of H so some of them is suitable to define the required fold out  $G_1$  of G.  $\Box$ 

## **3.1** When $ex(n, A) = \Theta(n^2)$

Here we collect all cases of  $\mathcal{A} \subset \mathcal{F}$  when we can prove that  $ex(n, \mathcal{A}) = \Theta(n^2)$ .

**Proposition 4.** Assume that  $\mathcal{A} \subset \mathcal{F} \setminus F_2$  and  $\mathcal{A} \cap \{F_1, F_3, F_7\} \neq \emptyset$ . Then  $ex(n, \mathcal{A}) = \Theta(n^2)$ .

**Proof.** The first condition ensures that the members of  $\mathcal{A}$  cannot be pierced by one vertex, thus Construction (C3) shows that  $ex(n, \mathcal{A}) = \Omega(n^2)$ . On the other hand,  $F_7$  is a fold out of  $F_3$  and  $F_1$  is a fold out of  $F_7$  thus by Lemma 2 (and by the second condition of the proposition)

$$ex(n, F_1) \le ex(n, F_7) + O(n^2) \le ex(n, F_3) + O(n^2) \le \frac{7}{2} \binom{n}{2} + O(n^2)$$

where the upper bound of  $ex(n, F_3)$  is Füredi's result [4].  $\Box$ 

**Proposition 5.** Assume that  $\mathcal{A} \subset \mathcal{F} \setminus F_3$  and  $\mathcal{A} \cap \{F_1, F_7\} \neq \emptyset$ . Then  $ex(n, \mathcal{A}) = \Theta(n^2)$ .

**Proof.** To show that  $ex(n, F) = \Omega(n^2)$ , consider the Construction (C3), it contains only  $F_3$  from  $\mathcal{F}$ . The upper bound follows by the argument of Proposition 4.  $\Box$ 

**Proposition 6.** Assume that  $\{F_2, F_3\} \subset \mathcal{A} \subset \{F_2, F_3, F_4, F_5, F_7\}$ . Then  $ex(n, \mathcal{A}) = \Theta(n^2)$ .

**Proof.** To show that  $ex(n, \mathcal{A}) = \Omega(n^2)$ , consider

• (C4) Steiner triple systems without  $F_4$  (the Pasch configuration), they do not contain any member of  $\mathcal{A}$ .

The upper bound follows from [4] since  $F_3 \in \mathcal{A}$ .  $\Box$ 

**Proposition 7.** Assume that  $\{F_2, F_3, F_6\} \subset A \subset \{F_2, F_3, F_5, F_6, F_7\}$ . Then  $ex(n, A) = \Theta(n^2)$ .

**Proof.** To show that  $ex(n, A) = \Omega(n^2)$ , consider

• (C5) Steiner triple systems without  $F_6$  (projective Steiner triple systems), they do not contain any member of  $\mathcal{A}$ .

The upper bound follows again from [4] since  $F_3 \in \mathcal{A}$ .  $\Box$ 

Note that Proposition 4 covers 56 cases, Proposition 5 adds 24 further cases, Propositions 6, 7 add 8 plus 4 further cases. These 92 cases are the ones when  $ex(n, A) = \Theta(n^2)$  follows from known results.

### 3.2 Unsolved cases

The 18 unsolved cases are grouped as follows.

- 1.  $ex(n, \{\mathcal{A} \cup F_6\})$  where  $\mathcal{A} \subseteq \{F_2, F_4, F_5\}$  (8 cases)
- 2.  $ex(n, \{F_2, F_5\}), ex(n, F_5)$
- 3.  $ex(n, \{F_2, F_4, F_5\}), ex(n, \{F_4, F_5\})$
- 4.  $ex(n, \{F_2, F_4\}), ex(n, F_4)$
- 5.  $ex(n, \mathcal{A})$  where  $\{F_2, F_3, F_4, F_6\} \subseteq \mathcal{A} \subseteq \{F_2, F_3, F_4, F_5, F_6, F_7\}$  (4 cases)

The upper bounds for the unknown cases can be compared by using Lemma 2. For example, observing that  $F_6$  is a fold out of  $F_4$  and of  $F_5$ , moreover  $F_5$  is a fold out of  $F_2$ , Lemma 2 implies

**Proposition 8.** Let  $\mathcal{A}$  be any subset of  $\{F_2, F_4, F_5\}$ . Then

$$\operatorname{ex}(n, F_6) \le \operatorname{ex}(n, \{\mathcal{A} \cup F_6\}) \le \operatorname{ex}(n, F_6) + 7\binom{n}{2}.$$

A lower bound  $\Omega(n^2)$  for the first four groups of unknown cases can be obtained from Construction (C3). Lower bounds for the fifth group of unknown cases can be given by well studied functions introduced in [1]. Let ex(n, (6, 3)) be the maximum number of triples in a triple system that does not contain three triples inside any six vertices. Since all members of  $\mathcal{F}$  except  $F_1$  contain three triples inside six vertices an almost quadratic lower bound of Construction (C1) comes from [9] for the four unsolved cases in group 5. A quadratic upper bound is from [4] since  $F_3 \in \mathcal{A}$ . Thus we get

**Proposition 9.** If  $\{F_2, F_3, F_4, F_6\} \subseteq \mathcal{A} \subseteq \{F_2, F_3, F_4, F_5, F_6, F_7\}$  then

$$\exp(n, (6, 3)) \le \exp(n, \mathcal{A}) = O(n^2)$$

In fact, the lower bound of Proposition 9 can be changed to ex(n, (7, 4)) (the maximum number of triples in a triple system that does not contain four triples inside any seven vertices).

**Acknowledgement** The authors are indebted to an unknown referee for pointing out some inaccuracies in the manuscript.

The research of the first author was supported in part by the Hungarian National Research, Development and Innovation Office NKFIH grant KH-130371 and NKFI-133819.

## References

- W. G. Brown, P. Erdős, V. T. Sós, Some extremal problems on r-graphs, in New directions in the theory of graphs, Proc. 3rd Ann Arbor Conference on Graph Theory, Academic Press, New York, 1973, 55–63.
- [2] P. Erdős, On extremal problems of graphs and generalized graphs, *Israel Journal of Mathematics* 2 (1964) 183–190.

- [3] P. Erdős, D. J. Kleitman, On coloring graphs to maximize the proportion of of multicolored k-edges, Journal of Combinatorial Theory 5 (1968) 164–169.
- [4] Z. Füredi, Hypergraphs in which all disjoint pairs have distinct unions, Combinatorica 4 (1984) 161–168.
- [5] Z. Füredi, L. Ozkahya, On 3-uniform hypergraphs without a cycle of given length, Discrete Applied Mathematics 216 (2017), 582–588.
- [6] D. Gerbner, A. Methuku, M. Vizer, Asymptotics for the Turán number of Berge- $K_{2,t}$ , arXiv:1705.04134v2
- [7] D. Mubayi, Some exact results and new asymptotics for hypergraph Turán numbers, Combinatorics, Probability and Computing 11 (2002) 299–309.
- [8] H. Leffmann, K. T. Phelps, V. Rödl, Extremal problems for triple systems, *Journal of Combinatorial Designs* 1. (1993) 379–394.
- [9] I. Z. Ruzsa, E. Szemerédi, Triple systems with no six points carrying three triangles, in: Combinatorics, Vol. II. Coll. Math. Soc. J. Bolyai 18 North-Holland, 1978, 939–945.