

HOW MANY ATOMS CAN BE DEFINED BY BOXES?

A. GYÁRFÁS, J. LEHEL and ZS. TUZA

Received 3 May 1983

We study the function $b(n, d)$, the maximal number of atoms defined by n d -dimensional boxes, i.e. parallelepipeds in the d -dimensional Euclidean space with sides parallel to the coordinate axes.

We characterize extremal interval families defining $b(n, 1) = 2n - 1$ atoms and we show that $b(n, 2) = 2n^2 - 6n + 7$.

We prove that for every d ,

$$b^*(d) = \lim_{n \rightarrow \infty} b(n, d)/n^d \text{ exists and } 1 \leq (d/2) \sqrt[d]{b^*(d)} \leq e.$$

Moreover, we obtain $b^*(3) = 8/9$.

1. Introduction

Let $\mathcal{A}_n = \{A_1, A_2, \dots, A_n\}$ be a family of non-empty sets. The elements $x, y \in \bigcup_{i=1}^n A_i$ are said to be equivalent if for all i , $1 \leq i \leq n$, $x \in A_i$ if and only if $y \in A_i$.

The equivalence classes are called the *atoms* of \mathcal{A}_n .

R. Rado studied the number of atoms for families given by immersion in a Euclidean space ([5]). In particular, he raised the problem of determining the maximal number of atoms in a family of n d -dimensional boxes, i.e. n parallelepipeds of the d -dimensional Euclidean space \mathbf{R}^d , with sides parallel to the coordinate axes. This number is denoted by $b(n, d)$ and families defining $b(n, d)$ atoms are called *extremal*. R. Rado proved in [5] that $b(n, 1) = 2n - 1$ and gave upper bounds for $b(n, d)$ and a lower bound for $b(n, 2)$.

The problem of determining the number (or the maximal number) of atoms for a family of figures of the Euclidean space is apparently new. However, some of our results (e.g. Remark 4.6.) show that this problem closely relates to a more familiar area of combinatorial geometry: to find the number (or maximal number) of *connected regions* defined by a family of figures. An excellent review is given by B. Grünbaum in [3].

This paper presents results on $b(n, d)$. We use the well-known notion of the *overlap graph* (cf. [2]) to study the 1-dimensional case, i.e. the number of atoms defined by families of *intervals*. We show that the number of atoms defined by n intervals is $2n - c$ where c denotes the number of connected components in the overlap graph (Theorem 2.1). As a consequence, we get a characterization of extremal interval families. An interval family is extremal if and only if its overlap graph is connected (Corollary 2.2).

Rectangles or 2-dimensional boxes are discussed in section 3. We introduce the notion of *overlay index* and we show its role related to the number of atoms in rectangle families (Theorems 3.1 and 3.2). These investigations lead to the main result of section 3: $b(n, 2) = 2n^2 - 6n + 7$ (Theorem 3.5). We note that the extremal families have an interesting structure, their complete description is given in [4].

We do not know the exact value of $b(n, d)$ for $d \geq 3$. In section 4 we show the existence of

$$b^*(d) = \lim_{n \rightarrow \infty} b(n, d)/n^d$$

for every fixed d (Theorem 4.3). We prove that $b^*(3) = 8/9$ (Corollary 4.7). For $d \geq 4$ we show that for a suitable constant $c \geq 1$, $c \leq (d/2) \sqrt[d]{b^*(d)} \leq e$ (Corollary 4.4). (The best value of c , known for us, is $\sqrt[3]{3}$ if d is a multiple of 3.)

Throughout the paper we impose two restrictions on the family of boxes, having no effect on the maximal number of atoms. We assume that the boxes are *closed*. Applying small enlargements for the boxes of a family, the number of atoms does not decrease, therefore we always assume that the *boundary hyperplanes of the boxes are all different*.

We note that extremal families of intervals may contain disjoint intervals. However, for higher dimensions the situation changes: *every extremal system must have non-empty intersection if $d \geq 2$* (Lemma 3.3).

2. Intervals

Let $\mathcal{I} = \{I_1, I_2, \dots, I_n\}$ be a family of intervals (all closed and having no common endpoints). Let $a(\mathcal{I})$ denote the number of atoms defined by \mathcal{I} . The *overlap graph* of \mathcal{I} , $G(\mathcal{I})$, is defined on the vertex set $\{1, 2, \dots, n\}$ and uv is an edge of $G(\mathcal{I})$ if and only if I_u and I_v overlap, i.e., they intersect but neither properly contains the other. As far as we know, overlap graphs were used first by Fulkerson and Gross [1] in the study of interval families; for further results see [2].

Theorem 2.1. *If the overlap graph of an interval family \mathcal{I} consists of c connected components then \mathcal{I} defines $2|\mathcal{I}| - c$ atoms.*

Proof. Let G_1, G_2, \dots, G_c be the connected components of $G(\mathcal{I})$ and consider the subfamilies

$$\mathcal{I}_j = \{I_v \in \mathcal{I} : v \text{ is a vertex in } G_j\} \quad (1 \leq j \leq c).$$

Replace in \mathcal{I} the members of \mathcal{I}_1 by the interval $U = \bigcup_{I \in \mathcal{I}_1} I$: $\mathcal{I}' = (\mathcal{I} \setminus \mathcal{I}_1) \cup \{U\}$.

It is easy to see that U overlaps no interval of \mathcal{I}' and thus $a(\mathcal{I}) = a(\mathcal{I}_1) + a(\mathcal{I}')$.

The repeated application of this argument yields $a(\mathcal{I}) = \sum_{j=1}^c a(\mathcal{I}_j)$, consequently it is sufficient to show that if \mathcal{I} has a connected overlap graph then $a(\mathcal{I}) = 2|\mathcal{I}| - 1$.

The set $\bigcup_{I \in \mathcal{I}} I$ is an interval splitted by $2|\mathcal{I}| - 2$ endpoints into $2|\mathcal{I}| - 1$ open intervals. We will prove that points of different intervals belong to different atoms by showing that every atom defined by \mathcal{I} is convex. Indeed, let p and q be two arbitrary points of an atom A and denote by P the interval with endpoints p and q . Since P overlaps no interval of \mathcal{I} , the overlap graph of the subfamily $\mathcal{I}' = \{I \in \mathcal{I} : I \subset P\}$ and the overlap graph of the subfamily $\mathcal{I}'' = \{I \in \mathcal{I} : I \supset P\}$ belong to different components of $G(\mathcal{I})$. By the connectedness of $G(\mathcal{I})$ it follows that $\mathcal{I}' \neq \emptyset$ and $\mathcal{I}'' = \emptyset$, consequently $P \subset A$.

Corollary 2.2. *If \mathcal{I} is a family of intervals then*

- (i) $|\mathcal{I}| \cong a(\mathcal{I}) \cong 2|\mathcal{I}| - 1$;
- (ii) $a(\mathcal{I})$ can have every value between $|\mathcal{I}|$ and $2|\mathcal{I}| - 1$;
- (iii) $a(\mathcal{I}) = 2|\mathcal{I}| - 1$ if and only if $G(\mathcal{I})$ is connected. ■

Two extremal interval families are displayed on Fig. 1.




Fig. 1. Two extremal interval families with overlap graphs P_n and K_n

3. Rectangles

In this section we deal with families of 2-dimensional boxes, i.e. rectangles in the plane, and we determine $b(n, 2)$.

We introduce the notion of overlay index of families of rectangles as follows.

Let \mathcal{B} be a family of rectangles and v_1, v_2, v_3 and v_4 be the four vertices of some rectangle $R \in \mathcal{B}$. If the vertex v_i is contained by n_i rectangles different from R then $\omega(v_i) = \max \{0, n_i - 1\}$ is called the overlay index of v_i and

$$\omega(R) = \sum_{i=1}^4 \omega(v_i)$$

is called the *overlay index* of R . The overlay index of the whole family is defined as $\omega(\mathcal{B}) = \sum_{R \in \mathcal{B}} \omega(R)$ (cf. Fig. 2).

Theorem 3.1. *If \mathcal{B} is a family of n rectangles with non-empty intersection then \mathcal{B} defines at most $2n^2 - 5n + 5 - \omega(\mathcal{B})$ atoms.*




Fig. 2. Five rectangles with total overlay index 3

Proof. Suppose that the interior of the set $\bigcap_{B \in \mathcal{B}} B$ contains the coordinate origin. The boundaries of the rectangles split the plane into connected domains. Let \mathcal{D} be the set of domains which are covered by *at least two* rectangles of \mathcal{B} . Denote by \mathcal{D}_1 the set of those domains from \mathcal{D} which meet the positive orthant.

Every domain $D \in \mathcal{D}_1$ has an upper right corner which is either the intersection of the boundary lines of two rectangles $P, Q \in \mathcal{B}$ or it is the vertex of some $R \in \mathcal{B}$ and lies in some $S \in \mathcal{B}$. To get an upper bound for $|\mathcal{D}_1|$, we associate to \mathcal{D} the pair P and Q , or the pair R and S . In the second case there are $\omega_1(R)+1$ possibilities to choose S , where $\omega_1(R)$ denotes the overlay index of R at its vertex belonging to the positive orthant. Therefore $|\mathcal{D}_1| \leq \binom{n}{2} - \sum_{R \in \mathcal{B}} \omega_1(R)$ and since this argument is true for the other three orthants and for the corresponding sets $\mathcal{D}_2, \mathcal{D}_3$ and \mathcal{D}_4 , we get

$$(1) \quad \sum_{i=1}^4 |\mathcal{D}_i| \leq 4 \binom{n}{2} - \sum_{i=1}^4 \sum_{R \in \mathcal{B}} \omega_i(R) = 2n^2 - 2n - \omega(\mathcal{B}).$$

By taking into account the domains which meet several orthants, we give an upper bound for the cardinality of \mathcal{D} .

For $j=2, 3$ and 4 let $\mathcal{D}^{(j)} \subset \mathcal{D}$ be the set of domains which meet j orthants and intersect the coordinate cross $x \cup y$ in $j-1$ intervals. Denote by $\mathcal{D}^{(0)}$ the set consisting of the domain which contains the origin and r further twofold connected domains which meet $x \cup y$ in four intervals ($0 \leq r \leq n-2$). Each coordinate axis is splitted by the boundary segments of the domains of \mathcal{D} into $2n-3$ intervals, thus

$$|\mathcal{D}^{(2)}| + 2|\mathcal{D}^{(3)}| + 3|\mathcal{D}^{(4)}| + 4|\mathcal{D}^{(0)}| - 2 = 4n - 6,$$

and clearly $|\mathcal{D}^{(0)}| = r + 1$. From here and by inequality (1)

$$|\mathcal{D}| = \sum_{i=1}^4 |\mathcal{D}_i| - (|\mathcal{D}^{(2)}| + 2|\mathcal{D}^{(3)}| + 3|\mathcal{D}^{(4)}| + 3|\mathcal{D}^{(0)}|) \leq$$

$$\leq 2n^2 - 2n - \omega(\mathcal{B}) - (4n - 5 - r),$$

that is

$$(2) \quad |\mathcal{D}| \leq 2n^2 - 6n + 5 - \omega(\mathcal{B}) + r.$$

The atoms of \mathcal{B} not appearing in \mathcal{D} are those covered by just one rectangle. Obviously the number of these atoms is at most $n-r$ since the largest twofold connected domain of $\mathcal{D}^{(0)}$ surrounds at least r rectangles. Thus, using (2), we obtain: $a(\mathcal{B}) \cong |\mathcal{D}| + n - r \cong 2n^2 - 5n + 5 - \omega(\mathcal{B})$. ■

The projections of a rectangle family \mathcal{B} into the coordinate axes x and y give two interval families. Their overlap graphs, denoted by $G_h(\mathcal{B})$ and $G_v(\mathcal{B})$, are called the *horizontal* and *vertical overlap graphs* of \mathcal{B} .

Theorem 3.2. *Let \mathcal{B} be a family of rectangles with non-empty intersection. If the vertical and the horizontal overlap graphs of the rectangles are connected then the overlay index of \mathcal{B} is at least $|\mathcal{B}|-2$.*

Proof. Let $\mathcal{B} = \{B_1, B_2, \dots, B_n\}$ and $n \geq 3$. By the connectivity of the vertical overlap graph G_v , it can be assumed that the top-line of B_1 and the bottom-line of B_2 meet every rectangle B_i , $1 \leq i \leq n$. Moreover, one can suppose that the rectangles are indexed in such a way that for every j , $3 \leq j \leq n$,

- (a) the top-line of B_j or the bottom-line of B_j meets every rectangle B_k with $j < k \leq n$, and
- (b) for some index $i < j$, B_i and B_j are vertically overlapping: ij is an edge of G_v .

Indeed, supposing that the first $j-1$ rectangles of \mathcal{B} satisfy (a) and (b), the j 'th rectangle can be chosen as follows. There is a rectangle from $\mathcal{B} \setminus \{B_1, \dots, B_{j-1}\}$ which vertically overlaps some B_i , $1 \leq i \leq j-1$, since G_v is connected. If the top-line of B_j meets every rectangle from $\mathcal{B} \setminus \{B_1, \dots, B_i\}$ then let B_j be the rectangle whose bottom-line intersects every rectangle of $\mathcal{B} \setminus \{B_1, \dots, B_{j-1}\}$; otherwise, B_j will be the rectangle whose top-line meets every member of $\mathcal{B} \setminus \{B_1, \dots, B_{j-1}\}$.

Observe that among the eight vertices of B_1 and B_2 exactly two, say s and t belong to $B_1 \cap B_2$. Hence, every rectangle B_i , satisfying $3 \leq i \leq n$ and $\{s, t\} \cap B_i \neq \emptyset$, increases the overlay index of $\{B_1, B_2\}$, i.e. if $K = \{i: 3 \leq i \leq n, \{s, t\} \cap B_i = \emptyset\}$ then $\omega(B_1) + \omega(B_2) \cong n - 2 - |K|$. Thus, it is sufficient to show that

$$\sum_{3 \leq i \leq n} \omega(B_i) \cong |K|.$$

Let us consider a spanning tree T_h of the connected horizontal overlap graph and direct the edges of T_h in such a way that every vertex different from 1 have out-degree one. Denote by j^+ the endpoint of the edge starting from vertex j , $2 \leq j \leq n$.

By property (b), for every $j \in K$ there exists a rectangle $B_{j'}$, $j' < j$ and $j' < j^+$, such that $B_{j'}$ vertically overlaps one of B_j and B_{j^+} . Using property (a), it is easy to verify that there is a vertex p among the twelve vertices of $B_{j'}$, B_j and B_{j^+} which is




Fig. 3. The triplet (j', j, j^+) defines overlay index increment at vertex p

covered twice by these rectangles (see the two essentially different situations on Fig. 3.). If this p is a vertex of B_k ($k=j', j$ or j^+) then we say that the triplet (j', j, j^+) belongs to B_k and overlays p . Remark that no triplet belongs to B_1 or B_2 , since if $j \in K$ then B_j does not overlay the vertices of B_1 and B_2 .

Let p be the vertex of some rectangle B_k , $3 \leq k \leq n$, and denote by $E(p)$ the set of ordered pairs jj^+ defined by the triplets (j', j, j^+) belonging to B_k and overlaying p . Since ordered pairs defined by different triplets are different,

$$(3) \quad \sum_p |E(p)| \cong |\{(j', j, j^+): j \in K\}| = |K|.$$

If $E(p) \neq \emptyset$ for some vertex p of a given B_k then $E(p)$ is the edge set of a subforest of the tree T_h ; denote by $V(p)$ its vertex set. For the minimal element j_0 (or j_0^+) of $V(p)$, clearly $j_0 \notin V(p)$ holds, therefore $\omega(p) \cong |(V(p) \cup \{j_0\}) \setminus \{k\}| - 1 = |V(p)| - 1 \cong |E(p)|$. From here, by (3), $\sum_{3 \leq k \leq n} \omega(B_k) \cong \sum_p |E(p)| \cong |K|$ follows. ■

Lemma 3.3. *Every extremal family of d -dimensional boxes has non-empty intersection if $d \geq 2$.*

Proof. Let \mathcal{B} be an extremal family of n boxes in \mathbf{R}^d and suppose that there are disjoint intervals on some coordinate axis among the projections of the boxes of \mathcal{B} . We define an operation on \mathcal{B} which reduces the number of non-intersecting pairs of intervals on the coordinate axes in such a way that the number of box atoms does not decrease.

Consider the projection of \mathcal{B} on the i 'th coordinate axis and choose two disjoint intervals $[p_1, p_2]$ (the projection of $P \in \mathcal{B}$) and $[q_1, q_2]$ (the projection of $Q \in \mathcal{B}$) such that $p_2 < q_1$, moreover, no interval endpoints lie in the open segment (p_2, q_1) .

Change P by P' and Q by Q' , replacing $[p_1, p_2]$ by $[p_1, q_1]$ and $[q_1, q_2]$ by $[p_2, q_2]$. Then every atom defined by \mathcal{B} which has a representative point outside of the strip $S = \{(x_1, \dots, x_d) \in \mathbf{R}^d: p_2 \leq x_i \leq q_1\}$ remains atom for the modified family \mathcal{B}' . If an atom A defined by \mathcal{B} is inside of S , i.e. it has no representative point outside of S , then for every $(a_1, \dots, a_d) \in A$, $(a_1, \dots, a_{i-1}, p_2, a_{i+1}, \dots, a_d) \in P$ and $(a_1, \dots, a_{i-1}, q_1, a_{i+1}, \dots, a_d) \in Q$. Thus different atoms inside the strip S remain different for \mathcal{B}' and since the set $P' \cap Q' \subset S$ contains them, they are all different from the atoms represented outside of S . Therefore $a(\mathcal{B}') \cong a(\mathcal{B})$, moreover if $P' \cap Q' \neq \emptyset$ and S is at least 2-dimensional then the atoms of \mathcal{B} inside of S can not cover $P' \cap Q'$ therefore $a(\mathcal{B}') > a(\mathcal{B})$ (cf. Fig. 4.).




Fig. 4. The boxes P and Q become intersecting and give birth to the black atom

After repeated applications of the above operation $P' \cap Q' \neq \emptyset$ occurs which would increase $a(\mathcal{B})$, contradicting the extremal property of \mathcal{B} . ■

Lemma 3.4. *For every n there exists an n -element extremal family of rectangles with connected vertical and horizontal overlap graphs.*

Proof. Let \mathcal{B} be an extremal family of n rectangles. By Lemma 3.3, $\bigcap_{B \in \mathcal{B}} B \neq \emptyset$.

Suppose that one of the overlap graphs of \mathcal{B} , say the vertical overlap graph G_v is not connected. This means that the top-line e of $P \in \mathcal{B}$ and a bottom-line f of $Q \in \mathcal{B}$ separate the rectangles of \mathcal{B} , i.e. $\{B \in \mathcal{B} : B \cap e \neq \emptyset\} = \{B \in \mathcal{B} : B \cap f \neq \emptyset\} \neq \mathcal{B}$.

To prove the lemma we show a transformation of \mathcal{B} which reduces the number of the connected components of G_v in such a way that the number of atoms does not decrease.

Let g be the closest top-line to e among the top-lines under e and suppose that g belongs to $R \in \mathcal{B}$. If we modify P and R exchanging their top-lines e and g then they become vertically overlapping (cf. Fig. 5.). On the other hand, the number of atoms does not decrease, since every atom meeting the open strip between e and g before the transformation has a representative point on the bottom-line f . ■




Fig. 5. Exchange of two top-lines

Theorem 3.5. *For every $n \geq 2$, $b(n, 2) = 2n^2 - 6n + 7$.*

Proof. Let \mathcal{B} be an extremal family of n rectangles. By Lemmas 3.3 and 3.4 one can suppose that the rectangles have a common point and both overlap graphs are connected. By Theorem 3.2, $\omega(\mathcal{B}) \geq n - 2$ and from Theorem 3.1, $b(n, 2) = a(\mathcal{B}) \geq 2n^2 - 5n + 5 - \omega(\mathcal{B}) \geq 2n^2 - 6n + 7$.

The next construction shows that $b(n, 2) \geq 2n^2 - 6n + 7$ for every $n \geq 2$. We give the rectangles R_k ($1 \leq k \leq n$) as direct products of closed intervals in a coordinate system of the plane:

$$R_k = [-k, k] \times [-n+1+k, n-1-k] \quad \text{if } 1 \leq k \leq n-2,$$

$$R_{n-1} = [-n+1, n-1] \times [-n, 0] \quad \text{and}$$

$$R_n = [-n, 0] \times [-n+1, n-1].$$

The family $\mathcal{R} = \{R_1, R_2, \dots, R_n\}$ splits the plane into $4 \binom{n-1}{2}$ unit squares and 3 further finite connected domains (cf. Fig. 6). All of these domains represent different atoms and therefore

$$b(n, 2) \cong a(\mathcal{R}) = 4 \binom{n-1}{2} + 3 = 2n^2 - 6n + 7.$$




Fig. 6. An extremal rectangle family

4. General estimations

When examining $b(n, d)$, it is useful to consider set systems consisting of so called corners. A d -dimensional corner at $\mathbf{v} = (v_1, \dots, v_d) \in \mathbf{R}^d$ is the set of points $\mathbf{x} = (x_1, \dots, x_d)$ such that $x_i \leq v_i$ whenever $1 \leq i \leq d$; in other words, corners are the translates of the negative orthant of \mathbf{R}^d .

We define $c(n, d)$ as the maximal number of atoms in a family consisting of n d -dimensional corners. We assume, just as in the case of boxes, that the corners have no common boundary hyperplanes. The close connection between $b(n, d)$ and $c(n, d)$ will be pointed out in Theorem 4.2 saying that their ratio is approximately 2^d .

First we give estimations on $c(n, d)$. It is straightforward that $c(n, 1) = n$ and $c(n, 2) = n + \binom{n}{2} = \binom{n+1}{2}$.

The key notion of this section is the *representation of atoms by index sets*. Consider a family of corners C_i at $\mathbf{v}_i = (v_1^i, \dots, v_d^i)$. Let A be an atom of this family and $I = \{i: A \subset C_i\}$. For $1 \leq k \leq d$, choose $i_k \in I$ such that C_{i_k} is minimal in the k 'th coordinate, i.e. $v_k^{i_k} < v_k^i$ if $i_k \neq j \in I$. We say that $I(A) = \{i_k: 1 \leq k \leq d\}$ represents A . Of course, $|I(A)| \leq d$ and different atoms are represented by different index sets.

Theorem 4.1. For every fixed $d \geq 3$,

$$\binom{n}{d} - O(n^{d-1}) \leq c(n, d) \leq \frac{2}{d} \binom{n}{d} + O(n^{d-1}).$$

Proof. 1. Upper bound. Let us consider a family of corners C_i at $\mathbf{v}_i = (v_1^i, v_2^i, \dots, v_d^i)$, $i = 1, \dots, n$. The number of atoms A with $|I(A)| < d$ is at most

$$(4) \quad \binom{n}{1} + \dots + \binom{n}{d-1} = O(n^{d-1}).$$

Let $c_0(n, d)$ denote the maximal number of atoms A with $|I(A)| = d$ in families of n d -dimensional corners and assume that $c_0(n, d)$ is achieved by our family C_1, \dots, C_n . According to (4),

$$(5) \quad c(n, d) \cong c_0(n, d) + \sum_{i=1}^{d-1} \binom{n}{i}.$$

We assume that the corners are indexed in such a way that for every k , $1 \leq k \leq d$, the corner vertices satisfy $\min_{k \leq i \leq n} v_k^i = v_k^k$. Any $(d-1)$ -element subfamily of $\{C_{d+1}, \dots, C_n\}$ has a member C_m such that, in this subfamily, \mathbf{v}^m is minimal in the j_1 'th and the j_2 'th coordinates ($1 \leq j_1 < j_2 \leq d$). Let I be the index set of this subfamily completed by some index k , $1 \leq k \leq d$, different from j_1 and j_2 . Then I does not represent any atom, since one of \mathbf{v}^m and \mathbf{v}^k gives the minimum in two different coordinates. The d -element subsets with the properties above can be chosen in $(d-2) \binom{n-d}{d-1}$ different ways. Repeating this procedure $[n/d]$ times for the subfamilies $\{C_{d+1}, \dots, C_n\}$, $\{C_{2d+1}, \dots, C_n\}$, and so on, we obtain

$$(6) \quad c_0(n, d) \cong \binom{n}{d} - (d-2) \sum_{i=1}^{[n/d]} \binom{n-id}{d-1}.$$

Since $(d-2) \sum_{i=1}^{[n/d]} \binom{n-id}{d-1} = \frac{d-2}{d} \binom{n}{d} - O(n^{d-1})$, (4), (5) and (6) imply $c(n, d) \cong \frac{2}{d} \binom{n}{d} + O(n^{d-1})$.

2. Lower bound. We define a family of n corners at vertices $\mathbf{v}^1, \dots, \mathbf{v}^n$ in the following way:

$$v_j^i = \begin{cases} i & \text{if } i \equiv j \pmod{d}, \\ 2n-i & \text{otherwise} \end{cases}$$

Every d -element set $I \subset \{1, \dots, n\}$ containing pairwise non-congruent indices mod d represents some atom for the family, since, for every $k \in I$, \mathbf{v}^k is minimal in the j 'th coordinate where $k \equiv j \pmod{d}$, $1 \leq j \leq d$. Consequently, $c(n, d) \cong (n/d)^d$ for every n divisible by d .

For every n , $(n/d)^d - [n/d]^d = O(n^{d-1})$, therefore $c(n, d) \cong (n/d)^d - O(n^{d-1})$ follows. ■

Theorem 4.2. For every $d \geq 1$, $\lim_{n \rightarrow \infty} \frac{b(n, d)}{c(n, d)} = 2^d$.

Proof. We proceed in 3 steps, proving:

$$(a) \quad b(n, d) \leq 2^d c(n, d)$$

$$(b) \quad b(n, d) \geq 2^d c(n-d, d)$$

$$(c) \quad \lim_{n \rightarrow \infty} \frac{c(n-d, d)}{c(n, d)} = 1.$$

(a) Let the boxes B_1, \dots, B_n form a family having $b(n, d)$ atoms. By Lemma 3.3, we suppose that the origin is inside $B_1 \cap \dots \cap B_n$. Then, in an arbitrary orthant of \mathbf{R}^d , the boxes can be considered as corners, therefore in a fixed orthant they can define at most $c(n, d)$ atoms. As the number of orthants is exactly 2^d , $b(n, d) \leq 2^d c(n, d)$ follows.

(b) Consider an extremal family of $n-d$ corners in \mathbf{R}^d , and suppose that the origin is contained by every corner. Let C_i be the corner at $\mathbf{v}^i = (v_1^i, \dots, v_d^i)$, $d+1 \leq i \leq n$. Then the points $\{(y_1, \dots, y_d) : |y_1| = v_1^i, \dots, |y_d| = v_d^i\}$ can be taken as the vertices of a box B_i . Let $m > \max\{v_j^i : 1 \leq j \leq d, d+1 \leq i \leq n\}$ and for every k , $1 \leq k \leq d$, $B_k = \{(x_1, \dots, x_d) : -m \leq x_i \leq m, x_k \leq 0\}$. The definition of B_1, \dots, B_d ensures that the $2^d c(n-d, d)$ atoms defined by B_{d+1}, \dots, B_n in the orthants are different.

(c) Every index $i \leq n$ is contained by at most $\binom{n-1}{d-1} + o(n^{d-1})$ representing sets $I(A) \subset \{1, \dots, n\}$. Therefore, deleting d arbitrary corners from an extremal family with $c(n, d)$ atoms, we have

$$c(n, d) \leq c(n-d, d) + d \binom{n-1}{d-1} + o(n^{d-1}) = c(n-d, d) + o(n^d).$$

By Theorem 4.1, $o(n^d)$ can be replaced by $o(c(n, d))$ and the proof is done because, on the other hand, $c(n, d) \geq c(n-d, d)$ is trivial. ■

Theorem 4.3. For every $d \geq 1$, $b^*(d) = \lim_{n \rightarrow \infty} \frac{b(n, d)}{n^d}$ exists.

Proof. By Theorem 4.2, it is enough to prove that

$$c^*(d) = \lim_{n \rightarrow \infty} \frac{c(n, d)}{n^d}$$

exists. To achieve this, we show the existence of

$$(7) \quad \lim_{n \rightarrow \infty} \frac{\binom{n}{d} - c_0(n, d)}{n^d};$$

with the $c_0(n, d)$ defined in the proof of Theorem 4.1 as the maximal number of atoms represented by d -element index sets in a family of n d -dimensional corners. From (5) $c(n, d) - c_0(n, d) = o(n^d)$.

The proof is based on the following general observation. Let \mathcal{H} be a class of d -uniform hypergraphs without multiple edges. Suppose that \mathcal{H} is closed under

the operation of vertex deletion and, for every n , there exists an $H \in \mathcal{H}$ on n vertices. Define $f(n) = \min \{|E(H)| : H \in \mathcal{H}, |V(H)| = n\}$, where $E(H)$ and $V(H)$ denote the set of edges and vertices of H respectively. Then $\lim_{n \rightarrow \infty} \frac{f(n)}{n^d}$ exists.

We apply this observation for the following class \mathcal{H} of hypergraphs. For every family of d -dimensional corners C_1, \dots, C_n , a d -uniform hypergraph H is defined: $V(H) = \{1, \dots, n\}$ and a d -element set I is an edge of H if and only if I does not represent an atom in the family. Now clearly $f(n) = \binom{n}{d} - c_0(n, d)$, therefore the existence of (7) follows. ■

Corollary 4.4. For every $d \geq 1$, $1 \leq \frac{d}{2} \sqrt[d]{b^*(d)} \leq e$. ■

The methods presented here lead to sharp results in the 3-dimensional case.

Theorem 4.5. For every $n \geq 1$, $c(n, 3) = \left\lfloor \frac{(n+1)^3 + 1}{9} \right\rfloor$.

Proof. For $d=3$, (5) and (6) imply that

$$(8) \quad c(n, 3) \leq n + \binom{n}{2} + \binom{n}{3} - \sum_{i=1}^{\lfloor n/3 \rfloor} \binom{n-3i}{2}.$$

Let \mathcal{C} be the family of n corners given in the proof for the lower bound of Theorem 4.1. Since no corner contains any other corner of \mathcal{C} , every one- and two-element index set represents an atom.

Let $\{k, l, m\}$ be a three-element index set with $1 \leq k < l < m \leq n$. Obviously, this set does not represent an atom of \mathcal{C} if and only if $k \equiv l \pmod{3}$. The number of these sets with $3(i-1) + 1 \leq k \leq 3i$ is $\binom{n-3i}{2}$ for every i , $1 \leq i \leq \lfloor n/3 \rfloor$. Therefore

$$(9) \quad c(n, 3) \geq n + \binom{n}{2} + \binom{n}{3} - \sum_{i=1}^{\lfloor n/3 \rfloor} \binom{n-3i}{2}.$$

The explicit form of the right hand side of (9) is $\left\lfloor \frac{(n+1)^3 + 1}{9} \right\rfloor$, thus by (8) and (9) the theorem follows. ■

Remark 4.6. The atoms defined by families of corners are connected regions therefore $c(n, d)$ gives the maximal number of connected regions defined by n corners. Theorem 4.5, for example can be stated as follows: n translates of an orthant in \mathbb{R}^3 divides \mathbb{R}^3 into at most $\left\lfloor \frac{(n+1)^3 + 1}{9} \right\rfloor + 1$ connected regions.

Corollary 4.7. $b^*(3) = 8/9$. ■

Conjecture 4.8. For $n \geq 3$, $b(n, 3) = 8 \left\lfloor \frac{(n-2)^3 + 1}{9} \right\rfloor + 7$.

References

- [1] D. R. FULKERSON and O. A. GROSS, Incidence matrices with the consecutive 1's property, *Bull. Amer. Math. Soc.* **70** (1964), 681—684.
- [2] M. C. GOLUBIC, *Algorithmic Graph Theory and Perfect Graphs*, Academic Press, 1980.
- [3] B. GRÜNBAUM, *Arrangements and spreads*, Amer. Math. Soc., Providence, 1972.
- [4] A. GYÁRFÁS, J. LEHEL and Zs. TUZA, The structure of rectangle families dividing the plane into maximal number of atoms, *Discrete Mathematics* **52** (1984), 177—198.
- [5] R. RADO, Atoms of families of sets, *Combinatorica* **2** (1982), 311—314.

András Gyárfás, Jenő Lehel, Zsolt Tuza

*Computer and Automation Institute of the
Hungarian Academy of Sciences
H-1111 Budapest, Kende u. 13—17.*

Hungary