

Hamiltonian cycles in the generating graphs of finite groups

Attila Maróti

(Rényi Institute of Mathematics, Budapest, Hungary)

Bath, August, 2009

What is the talk about?

This talk is based on the following draft.

Authors:

- ▶ Thomas Breuer,
- ▶ Robert M. Guralnick,
- ▶ Andrea Lucchini,
- ▶ A. M.,
- ▶ Gábor P. Nagy.

Title: Hamiltonian cycles in the generating graphs of finite groups.

The generating graph

Definition (Notation)

Let G be a non-trivial finite group.

Let $\Gamma(G)$ be the (simple) graph defined on the non-trivial elements of G in such a way that two distinct vertices are connected by an edge if and only if they generate G .

Why is the generating graph interesting?

1. Solvable groups

The **clique number** of a finite simple graph is the size of a largest complete subgraph of the graph. The **chromatic number** of a finite simple graph is the smallest number of colors needed to color the vertices of the graph in such a way that the endpoints of every edge receive different colors.

Theorem (Lucchini, M, 2008).

Let G be a finite group of Fitting height at most 2. Then the clique number and the chromatic number of $\Gamma(G)$ are equal.

Conjecture (Lucchini, M, 2008).

Let G be a finite solvable group. Then the clique number and the chromatic number of $\Gamma(G)$ are equal.

Why is the generating graph interesting?

2. Finite simple groups

Theorem (Steinberg, 1962; Aschbacher, Guralnick, 1984).

Every finite simple group can be generated by 2 elements.

What can be said about $\Gamma(G)$ when G is a finite simple group?

In the literature, many theorems about **finite simple groups** can be phrased in terms of the generating graph.

Theorem (Guralnick, Kantor, 2000).

If G is a finite simple group of order larger than 2, then there is no isolated vertex in $\Gamma(G)$.

Theorem (Guralnick, Shalev, 2003).

For every sufficiently large finite simple group G the diameter of the graph $\Gamma(G)$ is at most 2.

Theorem (Breuer, Guralnick, Kantor, 2008).

If G is a finite simple group of order larger than 2, then the diameter of the graph $\Gamma(G)$ is at most 2.

The conjecture of Breuer, Guralnick, and Kantor

The previous three theorems point in one direction...

Conjecture (Breuer, Guralnick, Kantor, 2008).

Let G be a finite group of order larger than 2. Then the following three conditions are equivalent.

1. G/N is cyclic for every non-trivial normal subgroup N of G ;
2. there is no isolated vertex in $\Gamma(G)$;
3. the diameter of $\Gamma(G)$ is at most 2.

Burness, Guest, and Guralnick are working on this conjecture. They have reduced the problem to the case where G is **almost simple**.

For which finite groups G does
there exist a Hamiltonian cycle in $\Gamma(G)$?

(A Hamiltonian cycle in a graph is a cycle that visits each vertex exactly once.)

Solvable groups

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

Let G be a finite **solvable** group of order larger than 3. Then the following four conditions are equivalent.

1. G/N is cyclic for every non-trivial normal subgroup N of G ;
2. there is no isolated vertex in $\Gamma(G)$;
3. the diameter of $\Gamma(G)$ is at most 2;
4. there is a Hamiltonian cycle in $\Gamma(G)$.

Conjecture.

We may remove the word **solvable** from the statement of the above theorem. (The implications 4. implies 2. implies 1. are trivial.)

Pósa's criterion

For solvable groups G it is somewhat easy to exhibit a Hamiltonian cycle in $\Gamma(G)$ (provided that a Hamiltonian cycle exists). But this is not always the case for finite groups in general.

Theorem (Pósa).

Let Γ be a simple graph on m vertices whose sequence of vertex degrees is $d_1 \leq \dots \leq d_m$. If $d_k \geq k + 1$ for every positive integer k less than $m/2$, then there exists a Hamiltonian cycle in Γ .

Finite simple groups

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

For every sufficiently large finite simple group G the graph $\Gamma(G)$ satisfies Pósa's criterion hence there exists a Hamiltonian cycle in $\Gamma(G)$.

The proof relies on the following papers:

- ▶ Babai-Hayes (2006);
- ▶ Fulman-Guralnick (2008);
- ▶ Guralnick-Kantor (2000);
- ▶ Liebeck-Shalev (1996).

Chvátal's criterion

If G is a finite group having a subgroup of index 2, then there is little chance that the graph $\Gamma(G)$ satisfies Pósa's criterion. In this case we use (a weaker form of) Chvátal's criterion.

Theorem (weaker form) (Chvátal).

Let Γ be a graph on m points with sequence of vertex degrees $d_1 \leq \dots \leq d_m$. Suppose that we have $m - k \leq d_{m-k}$ for all positive integers k with $k < m/2$. Then there exists a Hamiltonian cycle in Γ .

The Bondy-Chvátal closure

Pósa's and Chvátal's criteria are often combined with the following combinatorial tool.

Definition

Let Γ be a simple graph on m points. Let the degree of a vertex v be denoted by $d(v)$. The **Bondy-Chvátal closure** of the graph Γ is the graph $c/(\Gamma)$ obtained from Γ in such a way that there is an edge between vertices u and v if and only if there is an edge between u and v in Γ or there is no edge between u and v in Γ but $d(u) + d(v) \geq m$.

Theorem (Bondy, Chvátal, 1976).

The graph Γ contains a Hamiltonian cycle if and only if the Bondy-Chvátal closure $c/(\Gamma)$ contains a Hamiltonian cycle.

Symmetric groups

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

For every sufficiently large symmetric group S_n the graph $cl(cl(cl(\Gamma(S_n))))$ satisfies Chvátal's criterion hence there exists a Hamiltonian cycle in $\Gamma(S_n)$.

The proof relies on the following papers:

- ▶ Babai-Hayes (2006);
- ▶ Breuer-Guralnick-Kantor (2008).

Wreath products

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

Let n be a prime power. For every sufficiently large nonabelian finite simple group S (independent of n) the graph $cl(cl(\Gamma(S \wr C_n)))$ satisfies Pósa's criterion hence there exists a Hamiltonian cycle in $\Gamma(S \wr C_n)$.

The proof relies on the following papers:

- ▶ Babai-Hayes (2006);
- ▶ Fulman-Guralnick (2008);
- ▶ Guralnick-Kantor (2000).

Sporadic simple groups

The character-theoretic computational methods of the Breuer-Guralnick-Kantor (2008) paper helped obtain the following result.

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

Let G be an almost simple group whose socle is a sporadic simple group which is different from the Monster. Then the graph $cl(cl(\Gamma(G)))$ satisfies Pósa's criterion and hence there exists a Hamiltonian cycle in $\Gamma(G)$.

We intend to deal with the case when G is the Monster group.

Small groups

The following is a weaker form of a conjecture we have already mentioned.

Conjecture (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

Let G be a finite group of order larger than 3. Then $\Gamma(G)$ contains a Hamiltonian cycle if and only if G/N is cyclic for every non-trivial normal subgroup N of G .

Theorem (Breuer, Guralnick, Lucchini, M, Nagy, 2009).

This conjecture is true for

- ▶ all groups of orders at most 10^5 ;
- ▶ all almost simple groups of orders at most 10^6 ;
- ▶ all nonabelian finite simple groups of orders at most 10^7 .

Thank you for your attention!