

## Problems and results on the differences of consecutive primes.

By P. ERDŐS in Syracuse (U. S. A.).

Let  $p_1 < p_2 < \dots$  be the sequence of consecutive primes. Put  $d_n = p_{n+1} - p_n$ . The sequence  $d_n$  behaves extremely irregularly. It is well known that  $\overline{\lim} d_n = \infty$  (since the numbers  $n! + 2, n! + 3, \dots, n! + n$  are all composite). It has been conjectured that  $d_n = 2$  for infinitely many  $n$  (i. e. there are infinitely many prime twins). This conjecture seems extremely difficult. In fact not even  $\underline{\lim} d_n < \infty$ , or even  $\underline{\lim} \frac{d_n}{\log n} = 0$  has ever been proved. A few years ago I proved<sup>1)</sup> by using Brun's method that

$$(1) \quad \underline{\lim} \frac{d_n}{\log n} < 1.$$

$\underline{\lim} \frac{d_n}{\log n} \leq 1$  is an immediate consequence of the prime number theorem. WESTZYNTHIUS<sup>2)</sup> proved in the other direction that

$$(2) \quad \overline{\lim} \frac{d_n}{\log n} = \infty.$$

In fact he show that for infinitely many  $n$ ,

$$d_n > \log n \cdot \log \log \log n / \log \log \log \log n.$$

I proved<sup>3)</sup> using Brun's method that for infinitely many  $n$

$$(3) \quad d_n > c \frac{\log n \cdot \log \log n}{(\log \log \log n)^2}.$$

CHEN<sup>4)</sup> proved (3) very much simpler without using Brun's method,

<sup>1)</sup> *Duke Math. Journal*, Vol. 6 (1940), p. 438—441.

<sup>2)</sup> *Comm. Phys. Math. Soc. Sci. Fenn.*, Helsingfors, Vol. 5 (1931), No. 25. p. 1—37.

<sup>3)</sup> *Quarterly Journal of Math.*, Vol. 6 (1935), p. 124—128. In this paper one can find some more literature on the difference of consecutive primes.

<sup>4)</sup> *Schriften des Math. Seminars und des Instituts für angewandte Math. der Univ. Berlin*, 4 (1938), p. 35—55.

and RANKIN<sup>5)</sup> proved that

$$(4) \quad d_n > c \frac{\log n \cdot \log \log n \cdot \log \log \log \log n}{(\log \log \log n)^2}$$

In the present note I prove the following

Theorem:

$$(5) \quad \overline{\lim} \frac{\min(d_n, d_{n+1})}{\log n} = \infty.$$

In other words to every  $c$  there exist values of  $n$  satisfying the inequalities  $d_n > c \log n$ ,  $d_{n+1} > c \log n$ .

It can be conjectured that  $\overline{\lim} \left( \frac{\min(d_n, d_{n+1}, \dots, d_{n+k})}{\log n} \right) = \infty$  for every  $k$ , but I cannot prove this for  $k > 1$ .

It can also be conjectured that  $\underline{\lim} \frac{\max(d_n, d_{n+1})}{\log n} < 1$ , but I cannot prove this either.

Proof of the Theorem<sup>6)</sup>. Let  $n$  be a large integer,  $m = \varepsilon \cdot \log n$ , where  $\varepsilon$  is a small but fixed number,  $f(m)$  tends to infinity together with  $m$  and  $f(m) = o(\log m)^{1/\varepsilon}$ ,  $N = \prod_{p_i \leq m} p_i$ ,  $q_i$  denotes the primes  $\leq (\log m)^2$ ,  $r_i$  the primes of the interval  $[(\log m)^2, m^{1/100 \log \log m}]$ ,  $s_i$  the primes of the interval  $(m^{1/100 \log \log m}, \frac{m}{2})$ , and  $t_i$  the primes satisfying  $\frac{m}{2} \leq t_i \leq m$ .

Our aim will be to determine a residue class  $x \pmod{N}$  so that

$$(6) \quad (x+1, N) = 1 \text{ and } (x+k, N) \neq 1 \text{ for all } |k| \leq mf(m) \text{ and } k \neq +1.$$

Suppose we already determined an  $x$  satisfying (6). Then we complete the proof as follows: Consider the arithmetic progression  $(x+1) + dN$ ,  $d=1, \dots$ . Since  $(x+1, N) = 1$  it represents infinitely many primes, in fact by a theorem of LINNIK<sup>7)</sup> the least prime it represents does not exceed  $N^{c_1}$  where  $c_1$  is an absolute constant independent of  $N$ . Now by the prime number theorem, or by the more elementary results of TCHEBICHEFF, we have

$$N^{c_1} = \left( \prod_{p_i \leq m} p_i \right)^{c_1} < e^{2m c_1} = n^{2\varepsilon c_1} < n^{1/8}$$

for  $\varepsilon < \frac{1}{4c_1}$ , or there exists a prime  $p_j$  satisfying

$$(7) \quad p_j < n^{1/8}, \quad p_j = (x+1) + dN.$$

<sup>5)</sup> *Journal of the London Math. Soc.*, Vol. 13 (1938), p. 242–247. For further results on the difference of consecutive primes see P. ERDŐS and P. TURÁN, *Bull. Amer. Math. Soc.*, Vol. 54 (1948).

<sup>6)</sup> We use the method of CHEN.

<sup>7)</sup> On the least prime in an arithmetical progression, I. The basic theorem, *Math. Sbornik*, Vol. 15 (57), No 2, p. 139–178. II. The Deuring–Heilbronn phenomenon, *Math. Sbornik*, Vol. 15 (57), p. 347–368.

It follows from (6) that

$$(8) \quad p_{j+1} - p_j \geq mf(m), \quad p_j - p_{j-1} \geq mf(m).$$

Thus from (7) and (8)

$$(9) \quad \frac{p_{j+1} - p_j}{\log p_j} \geq \frac{mf(m)}{\log n} = \varepsilon f(m) \rightarrow \infty, \quad \frac{p_j - p_{j-1}}{\log p_j} \geq \frac{mf(m)}{\log n} = \varepsilon f(m) \rightarrow \infty,$$

which proves (5) and our Theorem is proved.

Now we only have to find an  $x$  satisfying (6). Put

$$(10) \quad x \equiv 0 \pmod{q_i}, \quad x \equiv 0 \pmod{s_i}.$$

Let  $|k| \leq mf(m)$ , have no factor among the  $q$ 's and  $s$ 's. Then we assert that  $k$  is either  $\pm 1$  or a prime  $> \frac{m}{2}$  or has all its prime factors among the  $r$ 's. For if not then  $k$  would be greater than the product of the least  $r$  and the least  $t$ , i. e.

$$k \geq \frac{m}{2} (\log m)^2 > mf(m); \quad (f(m) = o(\log m))$$

an evident contradiction.

Denote by  $u_1, u_2, \dots, u_\xi$  the integers  $\leq |mf(m)|$  all whose prime factors are  $r$ 's. We estimate  $\xi$  as follows: We split the  $u$ 's into two classes. In the first class are the  $u$ 's which have less than  $10 \cdot \log \log m$  different prime factors. The number of these  $u$ 's is clearly less than

$$(11) \quad (m^{1/100 \log \log m} \cdot \log m)^{10 \log \log m} < m^{2/5}$$

(since the number of integers of the form  $p^\alpha$ ,  $p^\alpha < mf(m)$ ,  $p < m^{1/100 \log \log m}$  is less than  $m^{1/100 \log \log m} \cdot \log m$ ).

For the  $u$ 's of the second class  $v(u) \geq 10 \cdot \log \log m$  ( $v(u)$  denotes the number of different prime factors of  $u$ ). Thus from

$$\sum 2^{v(u)} < 2 \sum_{b=1}^{mf(m)} 2^{v(b)} < cmf(m) \cdot \log m < m (\log m)^2$$

we obtain that the number of the  $u$ 's of the second class is less than

$$(12) \quad \frac{m (\log m)^2}{2^{10 \log \log m}} < \frac{m}{(\log m)^2}.$$

Hence finally from (11) and (12)

$$(13) \quad \xi = o\left(\frac{m}{\log m}\right).$$

Denote now by  $v_1, v_2, \dots, v_\eta$  the integers of absolute value  $\leq mf(m)$  which do not satisfy the congruence (10). Then the  $v$ 's are either  $-1$  or are  $u$ 's, or of the form  $\pm p$ ,  $\frac{m}{2} < p \leq mf(m)$ . Thus by (13) and the results

of TCHEBICHEFF about primes

$$(14) \quad \eta < c \frac{mf(m)}{\log m}.$$

Suppose we already determined for  $i < j$  a residue class  $\lambda^{(i)} \pmod{r_i}$  so that

$$(15) \quad x \equiv \lambda^{(i)} \pmod{r_i}, \quad \lambda^{(i)} \not\equiv -1, \quad i = 1, 2, \dots, (j-1).$$

Denote by  $v_1^{(j)}, \dots, v_{\eta_j}^{(j)}$  the  $v$ 's which do not satisfy any of the congruences (15). There clearly exists a residue class  $\pmod{r_j}$  which contains at least  $\eta_j/r_j$  of the  $v$ 's. Denote this residue class by  $\lambda_1^{(j)}$ . If  $\lambda_1^{(j)} \not\equiv -1 \pmod{r_j}$  we put

$$(16) \quad x \equiv \lambda_1^{(j)} \pmod{r_j}.$$

If on the other hand  $\lambda_1^{(j)} \equiv -1 \pmod{r_j}$  we distinguish two cases: In the first case the residue class  $\lambda_1^{(j)} \pmod{r_j}$  contains less than  $\frac{1}{2} \eta_j$  of the  $v^{(j)}$ 's.

Then there clearly exists a residue class  $\lambda_2^{(j)} \not\equiv \lambda_1^{(j)} \pmod{r_j}$  which contains more than  $\eta_j/2r_j$  of the  $v^{(j)}$ 's. Put for these  $r_j$ 's

$$(17) \quad x \equiv \lambda_2^{(j)} \pmod{r_j}.$$

We continue this operation for all the  $r$ 's and let us first assume that for every  $r_j$  either  $\lambda_1^{(j)} \not\equiv -1 \pmod{r_j}$  or that the first case occurs. Denote by  $V_1, V_2, \dots, V_\varrho$  the  $v$ 's which do not satisfy the congruences (16) and (17). Clearly

$$(18) \quad \varrho \leq \eta \Pi \left(1 - \frac{1}{2r_j}\right) < c \frac{mf(m) \log \log m}{(\log m)^{3/2}} = o\left(\frac{m}{\log m}\right)$$

since

$$\frac{c_1}{\sqrt{\log z}} < \prod_{p \leq z} \left(1 - \frac{1}{2p}\right) < \frac{c_2}{\sqrt{\log z}}.$$

Put now

$$(19) \quad x \equiv -V_i \pmod{t_i}, \quad 1 \leq i \leq \varrho,$$

where  $t_i$  is chosen so that  $V_i - 1 \not\equiv 0 \pmod{t_i}$  and the different  $V_i$  correspond different  $t_i$ . This is always possible since the number of prime factors of  $V_i - 1$  is less than  $c \log m$  and number of  $t$ 's equals  $\pi(2m) - \pi(m)$ , and we have by (18) and the results of TCHEBICHEFF

$$\pi(2m) - \pi(m) > c_1 \frac{m}{\log m} > \varrho + c \log m.$$

For the  $t$ 's not used in (19) we put

$$(20) \quad x \equiv 0 \pmod{t_i}.$$

The congruences (10), (16), (17) and (20) determine  $x \pmod{N}$  so that (6) is clearly satisfied, which proves our Theorem in case the second case never occurs.