# A sharpening of Tusnády's inequality

Jenő Reiczigel<sup>1</sup>

Lídia Rejtő<sup>2,3</sup>

Gábor Tusnády<sup>3</sup>

November 24, 2011

#### Abstract

Let  $\varepsilon_1, \ldots, \varepsilon_m$  be i.i.d. random variables with

$$P(\varepsilon_i = 1) = P(\varepsilon_i = -1) = 1/2,$$

and  $X_m = \sum_{i=1}^m \varepsilon_i$ . Let  $Y_m$  be a normal random variable with the same first two moments as that of  $X_m$ . There is a uniquely determined function  $\Psi_m$  such that the distribution of  $\Psi_m(Y_m)$  equals to the distribution of  $X_m$ . Tusnády's inequality states that

$$|\Psi_m(Y_m) - Y_m| \le \frac{Y_m^2}{m} + 1.$$

Here we propose a sharpened version of this inequality. AMS 2000 subject classification. Primary 62E17; secondary 62B15 Key words and phrases. Quantile transformation; normal approximation; binomial distribution; Tusnády's inequality

### 1 Conjecture

Let  $\varepsilon_1, \ldots, \varepsilon_m$  be i.i.d. random variables with

$$P(\varepsilon_i = 1) = P(\varepsilon_i = -1) = 1/2,$$

and  $X_m = \sum_{i=1}^m \varepsilon_i$ . Let  $Y_m$  be a normal random variable with the same first two moments as that of  $X_m$ . Using quantile transformation we can

<sup>&</sup>lt;sup>1</sup>Szent István University, Department of Biomathematics and Informatics, Faculty of Veterinary Science, Budapest, Hungary

<sup>&</sup>lt;sup>2</sup>University of Delaware, Statistics Program, FREC, CANR, Newark, Delaware, USA

<sup>&</sup>lt;sup>3</sup>Alfréd Rényi Mathematical Institute of the Hungarian Academy of Sciences, Budapest, Hungary

see that there is a uniquely determined function  $\Psi_m$  such that the distribution of  $\Psi_m(Y_m)$  equals to the distribution of  $X_m$ . The central limit theorem implies that the function  $\Psi_m$  is close to the identity for large m. A sharp inequality of Tusnády [12] raised certain interest in the literature ([1],[2],[3],[4],[5],[6],[7],[8],[9],[10],[11],[13]).

Let us define the function f on the interval (0, 1) as

$$f(x) = \sqrt{(1+x)\log(1+x) + (1-x)\log(1-x)},$$

set  $f(0) = 0, f(1) = \sqrt{\log(4)}$ . Let us put

$$x_{k,m} = \frac{k - \frac{m}{2}}{\frac{m}{2}}$$

for positive even integers m with k such that  $m/2 < k \leq m$ , and set

$$p_{k,m} = P(X_m \ge 2k - m) = 2^{-m} \sum_{i=k}^m \binom{m}{i}.$$

Let us define the function Q on the reals as

$$Q(x) = \frac{1}{\sqrt{2\pi}} \int_x^\infty e^{-u^2/2} du.$$

With those ingredients our conjecture states that

$$Q(\sqrt{m}f(x_{k,m})) < p_{k,m} < Q(\sqrt{m}f(x_{k-1,m}))$$

holds true for  $\frac{m}{2} < k \leq m$ . Or more sharply

$$2(k-1) - \frac{m}{2} + 0.8964 < mf^{-1}(Q^{-1}(p_{k,m})/\sqrt{m}) < 2(k-1) - \frac{m}{2} + 1.0000$$
(1)

holds true with pessimal parameters m = k = 10. It implies that Tusnády's inequality is sharpened to

$$\left|\Psi_m(Y_m) - mf^{-1}\left(\frac{Y_m}{m}\right)\right| < 1.1036.$$

#### 2 Generalization

For an arbitrary random variable X let us consider the function on reals

$$R(t) = Ee^{tX}$$

restricting ourselves for distributions having finite momentum generators. Next we define P'(t)

$$\psi(t) = \frac{R(t)}{R(t)},$$
  

$$\alpha(x) = t \quad \text{iff} \quad \psi(t) = x,$$
  

$$\phi(x) = R(\alpha(x)) \exp(-x\alpha(x))$$

The probability  $P(\sum_{i=1}^{m} X_i \ge mx)$  is approximately  $\rho(x)^{-m}$  if x > EX. The function  $\rho$  depends on the distribution of X, it is the Chernoff function of X. Let us denote the Chernoff function of the distribution F of X by  $\rho_F$ , and the corresponding function for standard normal by  $\rho_G$ . The quantile transformation between the partial sums of distribution F with Gaussian ones resemble us to the equation

$$\rho_F(x) = \rho_G(y)$$

having the property that it gives sharp values for any m. Perhaps the error term is bounded with a bound depending on the distribution of X. For the case symmetrical binomial distribution the error term might be as small as that the quantile curve jumps over its limiting function: it is the informal explanation of our conjecture.

## 3 Numerical Illustration

The function  $\Psi_m$  is shown in Figure 1. called "step" for m = 50 with a rescaling for random variables

$$\xi_m = \frac{X_m}{m}, \quad \eta_m = \frac{Y_m}{m}.$$

The function f is called "limit", for the sequence of step functions goes to f after rescaling. The conjecture comes from the observation that the limit function crosses all steps near to their middle. Let us introduce the blow up error term

$$\Delta_{k,m} = 10 \left( 2k - 1 - m \ f^{-1} \left( \frac{1}{\sqrt{m}} \ Q^{-1} \left( \sum_{i=k}^{m} \binom{m}{i} 2^{-m} \right) \right) \right),$$

for  $0 < k \leq m/2$ . In Figure 1. it is labelled as "Delta". With these notations (1) is equivalent with  $0 < \Delta_{k,m} < 1.036$ . These error terms are shown in Figure 2. for  $2 \leq m \leq 1000$ . Figure 2. prompts the conjecture that even these curves are convergent. We are a bit perplexed: even the inequality

 $0 < \Delta_{1,2} < 1.036$  means that Q(0.723359) < 0.25 < Q(0.6435214). How can we prove such an inequality theoretically?

ACKNOWLEDGEMENT: We thank to Peter Harremoës for pointing out a mistake in the earlier version of the paper.

# References

- [1] Bretagnolle, J. and Massart, P. (1989). Hungarian constructions from the nonasymptotic viewpoint, *The Annals of Probability* **17**, 239–256.
- [2] Castelle, N. (2009). Improvement of two Hungarian bivariate theorems, Manuscript
- [3] Carter, A. and Pollard, D. (2004). Tusnády's inequality revisited, The Annals of Statistics 32, 2731–2741.
- [4] Csörgő, M. (2007). A glimpse of the KMT (1975): approximation of empirical processes by Brownian bridges via quantiles, Acta Sci. Math. (Szeged) 73, 349–366
- [5] Csörgő, M. and Révész, P. (1981). Strong Approximations in Probability and Statistics, Academic Press, New York.
- [6] Dudley, R. M. (1999). Notes on empirical processes, Lecture notes for a course given at Aarhus Univ., August 24, 1999.
- [7] Dudley, R. M. (2008). On the quantile transformation for asymmetric binomial and hypergeometric distributions, Manuscript
- [8] Major, P. (1999). The approximation of the empirical distribution function, Technical report, Alfréd Rényi Mathematical Institute of the Hungarian Academy of Sciences. Notes available from http://www.renyi.hu/~major/probability/empir.html, August 3, 1999.
- [9] Mason, D. M. (2001). Notes on the KMT Brownian bridge approximation to the uniform empirical process, In Asymptotic Methods in Probability and Statistics with Applications, (N. Balakrishnan, I. A. Ibragimov and V. B. Nevzorov, eds.) 351–369. Birkhäuser, Boston.
- [10] Massart, P. (2002). Tusnády's lemma, 24 years later, Ann. Inst. H. Poincaré Probab. Statist. 38, 991–1007.

- [11] Pollard, D. (2002). A user's guide to measure theoretic probability, Cambridge University Press
- [12] Tusnády, G. (1977). *Study of statistical hypotheses*, Dissertation for Habilitation, Hungarian Academy of Sciences, Budapest.
- [13] Zhou, H. H. (2006). A note on quantile coupling inequalities and their applications, Manuscript

# 4 Appendix

```
R- program of Figures 1 and 2.
```

```
Q=function(p) -qnorm(p)
G=function(x) ((1+x)*log(1+x)+(1-x)*log(1-x))**0.5
Ginv=function(u) {
GG=function(x) G(x)-u
uniroot(GG, c(0,1), f.lower=-u, f.upper=log(4)^.5-u, tol=10^-100)
}
m=50; k=m/2
sum=0; divisor=2**m; bin=
xx=c(1:k+1); yy=c(1:k+1); zz=c(1:k+1);
for (i in 1:k-1){
sum=sum+bin
x=(m-2*i)/m
y=Q(sum/divisor)/(m**.5)
b=Ginv(y)$root
yy[i+1]=y; xx[i+1]=x
bin=(m-i)*bin/(i+1)
zz[i+1]=10*(m-2*i-1-m*b)
xx[k+1]=0; yy[k+1]=0; zz[k+1]=0
kerx=c(0,1.25); kery=c(0,1.15)
plot(kerx, kery, type="n",xlab="eta", ylab="xi",
main="Figure1. Quantile transform, its limit and blownup error, m=50")
for (i in 1:k){
bb=seq(from=yy[i+1], to=yy[i], by=0.01)
cc=bb*0+1; cc=cc*xx[i+1]
points(bb,cc,type="l", col="blue", lwd=2)}
cc=seq(from=0, to=0.999, by=0.001)
bb=((1+cc)*log(1+cc)+(1-cc)*log(1-cc))**0.5
points(bb,cc, type="l", col="red", lwd=2)
points(yy,zz, type="l", col="green", lwd=2)
legend(locator(1),c("Limit","Step","Delta"),
lty=c(1,1,1),
col=c("red","blue","green"))
```

```
kerx=c(0,1.25); kery=c(0,1.15)
plot(kerx, kery, type="n",xlab="eta", ylab="Delta",
     main="Figure 2. The blownup error")
for (k in 1:500){m=2*k;
sum=0; divisor=2**m; bin=1
yy=c(1:k+1); zz=c(1:k+1);
for (i in 1:k-1){
sum=sum+bin
y=Q(sum/divisor)/(m**.5)
b=Ginv(y)$root
yy[i+1]=y;
bin=(m-i)*bin/(i+1)
zz[i+1]=10*(m-2*i-1-m*b)
yy[k+1]=0; zz[k+1]=0
if (k<100) clr="red" else
if (k<200) clr="blue" else
if (k<300) clr="purple" else
if (k<400) clr="gray" else clr="green"
points(yy,zz, type="l", col=clr)}
legend(locator(1),c("0<m <= 200","200<m<=400","400<m<=600",</pre>
"600<m<=800","800<m<=1000"),
lty=c(1,1,1,1,1),
col=c("red","blue","purple","gray","green"))
```



Figure1. Quantile transform, its limit and blownup error, m=50



Figure 2. The blownup error