

A note on triangulations of sum sets

K. Böröczky Jr. B. Hoffman

May 31, 2013

Abstract

In \mathbb{R}^2 , for finite sets A and B , we write $A+B = \{a+b : a \in A, b \in B\}$. We write $\text{tr}(A)$ to denote the common number of triangles in any triangulation of the convex hull of A using the points of A as vertices. We consider the conjecture that $\text{tr}(A+B)^{\frac{1}{2}} \geq \text{tr}(A)^{\frac{1}{2}} + \text{tr}(B)^{\frac{1}{2}}$. If true, this conjecture would be a discrete, two-dimensional analogue to the Brunn-Minkowski inequality. We prove the conjecture in three special cases.

1 Introduction

In this paper, we write A, B to denote finite subsets of \mathbb{R}^d , and $|\cdot|$ to stand for their cardinality. For objects X_1, \dots, X_k in \mathbb{R}^2 , $[X_1, \dots, X_k]$ denotes their convex hull. Our starting point is two classical results. One is due to Freiman from the 1960's; namely,

$$|A+B| \geq |A| + |B| - 1, \tag{1}$$

with equality if and only if A and B are arithmetic progressions of the same difference. The other result, the Brunn-Minkowski inequality dates back to the 19th century. It says that if $X, Y \subset \mathbb{R}^d$ are compact sets then

$$\lambda(X+Y)^{\frac{1}{d}} \geq \lambda(X)^{\frac{1}{d}} + \lambda(Y)^{\frac{1}{d}}$$

where λ stand for the Lebesgue measure, and equality holds if X and Y are convex homothetic sets. This theorem has been successfully applied to estimating the size of a a sumset say by Ruzsa, Green, Tao. In turn various discrete analogues of the Brunn-Minkowski inequality have been established in papers by Bollobás-Leader, Gardner-Gronchi, Green-Tao and most recently by Grynkiewicz-Serra in the planar case. All these papers use the

method of compression, which changes a finite set into a set better suited for sumset estimates, but cannot control the convex hull. See G.A. Freiman [1] and [2] for the earlier history, and I.Z. Ruzsa [4] and T. Tao, V. Vu [5] for thorough surveys.

Unfortunately the known analogues are not as simple in their form as the original Brunn-Minkowski inequality. A formula due to Gardner and Gronchi says that if A is not contained in any affine subspace of \mathbb{R}^d then

$$|A + B| \geq (d!)^{-\frac{1}{d}}(|A| - d)^{\frac{1}{d}} + |B|^{\frac{1}{d}}.$$

In this paper, we discuss a more direct version of the Brunn-Minkowski inequality in the plane, which would improve Freiman's inequality if both A and B are two-dimensional.

In the planar case ($d = 2$), a recent conjecture by Matolcsi and Ruzsa [3] might point to the right version of the Brunn-Minkowski inequality. Let A be a finite non-collinear point set in \mathbb{R}^2 . We write $\text{tr}(A)$ to denote the common number of triangles in any triangulation of $[A]$ using the points of A as vertices. If b_A and i_A denote the number of points of A in $\partial[A]$ and $\text{int}[A]$, then the Euler formula yields

$$\text{tr}(A) = b_A + 2i_A - 2. \tag{2}$$

If Π is a polygon whose vertices are in \mathbb{Z}^2 , and $A = \mathbb{Z}^2 \cap \Pi$, then Pick's theorem says that

$$\text{tr}(A) = 2\lambda(\Pi).$$

Now the Ruzsa-Matolcsi conjecture proposes that if A and B in the plane are not collinear then

$$\text{tr}(A + B)^{\frac{1}{2}} \geq \text{tr}(A)^{\frac{1}{2}} + \text{tr}(B)^{\frac{1}{2}}. \tag{3}$$

We note that equality holds if for a polygon Π whose vertices are in \mathbb{Z}^2 , and integers $k, m \geq 1$, we have $A = \mathbb{Z}^2 \cap k\Pi$ and $B = \mathbb{Z}^2 \cap m\Pi$.

In this paper, we verify (3) in some special cases. To present our main idea we note that if $\alpha, \beta > 0$, then

$$(\alpha + \beta)^2 \leq 2(\alpha^2 + \beta^2), \tag{4}$$

with equality if and only if $\alpha = \beta$. Thus Conjecture (3) follows from

$$\text{tr}(A + B) \geq 2[\text{tr}(A) + \text{tr}(B)]. \tag{5}$$

This inequality does not hold in general. For example, let Π be a polygon whose vertices are in \mathbb{Z}^2 , and let $A = \mathbb{Z}^2 \cap k\Pi$ and $B = \mathbb{Z}^2 \cap m\Pi$ for integers

$k, m \geq 1$. If $k \neq m$, then we have equality in the Brunn-Minkowski theorem for $X = [A]$ and $Y = [B]$. Still, as we verify, (5) holds in several interesting cases.

The triangulation conjecture (3) can be written in the following form.

Conjecture 1 (Main conjecture) *If A and B are finite non-collinear sets \mathbb{R}^2 , then*

$$\sqrt{2i_{A+B} + b_{A+B} - 2} \geq \sqrt{2i_A + b_A - 2} + \sqrt{2i_B + b_B - 2}.$$

In turn, (5) is equivalent with

$$2i_{A+B} + b_{A+B} \geq 4i_A + 4i_B + 2b_A + 2b_B - 6. \quad (6)$$

2 Remarks on the boundary

In the following, we need the notion of exterior normal. A vector u is an exterior normal at x_0 to $[A]$, where $x_0 \in A$, if

$$u \cdot x_0 = \max\{u \cdot x : x \in A\}.$$

It immediately follows that only points in the boundary of $[A]$ will have nonzero exterior normals. It also follows that if $a + b$ is a boundary point of $[A + B]$ for $a \in A$ and $b \in B$, then an exterior unit normal u at $a + b$ to $[A + B]$ is an exterior unit normal at a to $[A]$, and at b to $[B]$. We conclude the following.

Lemma 2 *If A and B are finite non-collinear sets in \mathbb{R}^2 , and $a \in A$ and $b \in B$, then $a + b$ lies on the boundary of $[A + B]$ with nonzero exterior unit normal vector u if and only if u is an exterior normal to $[A]$ at a and to $[B]$ at b .*

For a unit vector u , and finite set A , define the collinear set of points

$$A_u = \{x \in A : u \cdot x = \max_{y \in A} u \cdot y\}.$$

Lemma 3 *For any finite non-collinear sets A and B in \mathbb{R}^2 , we have*

$$b_{A+B} \geq b_A + b_B,$$

with equality if and only if $|A_u| \geq 2$ and $|B_u| \geq 2$ for a unit vector u imply that A_u and B_u are arithmetic progressions of the same difference.

Proof: For a finite collinear set C , let $S(C) = |C| - 1$; namely, the number of segments the points of C divide the line into. Therefore if C and D are contained in parallel lines, then $S(C + D) \geq S(C) + S(D)$, with equality if and only if either $|C| = 1$, or $|D| = 1$, or C and D are arithmetic progressions of the same difference. Applying this observation to $C = A_u$ and $D = B_u$ for each unit vector that is an exterior normal to a side of $[A + B]$ yields the lemma. \square

3 Sums with unique representation for each point

In this section we consider the case where representation of points in $A + B$ is unique. We say that the representation is unique when for all $x \in A + B$, if $x = a_1 + b_1$ and $x = a_2 + b_2$, then $a_1 = a_2$ and $b_1 = b_2$.

Theorem 4 *If the representation of points in $A + B$ is unique, then Conjecture 1 holds.*

Proof: From the previous section, we see that whether $x = a + b \in A + B$ lies on the boundary of $[A + B]$ depends only on the exterior normals of $a \in A$ and $b \in B$. So applying any transformation to A or B that preserves $|A + B|$, $\text{tr}(A)$, $\text{tr}(B)$, and the exterior normals of A and B will also preserve $\text{tr}(A + B)$. Note that scalar multiplication by ϵ , where $\epsilon A = \{\epsilon a : a \in A\}$ satisfies the latter three conditions immediately. Since the representation of points in $A + B$ is unique, picking ϵ so that the representation of points in $\epsilon A + B$ is also unique will satisfy the first condition.

We pick ϵ small enough so that, for fixed $b \in B$, letting $\epsilon A + b = \{a + b : a \in \epsilon A\}$, for any $x \in \epsilon A + B$, if $x \in [\epsilon A + b]$, then $x \in \epsilon A + b$. Geometrically, this amounts to shrinking A to a degree such that $\epsilon A + B$ looks like a little copy of A placed at each point in B . It follows that the representation of points in $\epsilon A + B$ is unique, and hence $\text{tr}(\epsilon A + B) = \text{tr}(A + B)$.

Assume without loss of generality that $\text{tr}(A) = \text{tr}(\epsilon A) \geq \text{tr}(B)$. We begin to draw lines between points in $\epsilon A + B$ to form a partial triangulation, which can be extended to a triangulation of $\epsilon A + B$. For each $b \in B$, draw lines on $\epsilon A + b$ that form a triangulation of that set. Then, consider a triangulation T of B . For each $b_1, b_2 \in B$ that are connected by a line in T , consider $\epsilon A + b_1$ and $\epsilon A + b_2$. Pick a point $b_1^* \in \epsilon A + b_1$ that has exterior normal $b_2 - b_1$ in $[\epsilon A + b_1]$. Pick a point $b_2^* \in \epsilon A + b_2$ that has exterior normal $b_1 - b_2$ in $[\epsilon A + b_2]$. Now, in $\epsilon A + B$, draw a line between b_1^* and b_2^* .

Geometrically, we have mimicked a triangulation of A at each little copy of A , and a triangulation of B on a large scale, treating each little copy of A as a point in B . Letting $\text{ptr}(\epsilon A + B)$ denote the number of polygons enclosed in this partial triangulation, it follows:

$$\text{tr}(A + B) = \text{tr}(\epsilon A + B) \geq \text{ptr}(\epsilon A + B) = |B|\text{tr}(A) + \text{tr}(B). \quad (7)$$

Conjecture 1 then follows from

$$\sqrt{|B|\text{tr}(A) + \text{tr}(B)} \geq \sqrt{\text{tr}(A)} + \sqrt{\text{tr}(B)}. \quad (8)$$

Since $|B| \geq 3$ and $\text{tr}(A) \geq \text{tr}(B)$, $(|B| - 2)\text{tr}(A) \geq \text{tr}(B)$ holds, which in turn implies (8). \square

4 The case $i_A = i_B = 1$

We see that Lemma 3 yields that (6), and in turn Conjecture 1 would follow from

$$2i_{A+B} \geq 4i_A + 4i_B + b_A + b_B - 6, \quad (9)$$

which we have already noted does not always hold. However, in the remainder of this paper we show it holds for two special cases. The proof of the first case is simple:

Theorem 5 *When $i_A = i_B = 1$, Conjecture 1 holds.*

Proof: From Lemma 2, it follows that if $a \in A_{\text{int}} = \{a \in A : a \in \text{int}([A])\}$, then $a + B \subset (A + B)_{\text{int}}$. So by (1), since i_A and i_B are nonempty, $i_{A+B} \geq i_A + |B| - 1$, and similarly $i_{A+B} \geq i_B + |A| - 1$. Thus, since $|A| = i_A + b_A$ and $|B| = i_B + b_B$, we have

$$2i_{A+B} \geq 2i_A + 2i_B + b_A + b_B - 2. \quad (10)$$

In the case that $i_A = i_B = 1$, (9) follows. \square

5 The case $|A| = b_A$ and $|B| = b_B$

We now turn to the case $|A| = b_A$ and $|B| = b_B$, or in other words, both A and B lie on the boundary of their convex hulls. In this case, (9) becomes

$$2i_{A+B} \geq b_A + b_B - 6. \quad (11)$$

The bad news is that (11) does not always hold. Let

$$\tilde{A} = \{(0, 0), (1, 0), (0, 1)\} = \{(x, y) \in \mathbb{N}^2 : x + y \leq 1\};$$

$$\tilde{B} = \{(0, 0), (1, 0), (0, 1), (2, 0), (1, 1), (0, 2)\} = \{(x, y) \in \mathbb{N}^2 : x + y \leq 2\}.$$

Therefore $|\tilde{A}| = b_{\tilde{A}} = 3$, $|\tilde{B}| = b_{\tilde{B}} = 6$, and $\tilde{A} + \tilde{B} = \{(x, y) \in \mathbb{N}^2 : x + y \leq 3\}$ yields $i_{\tilde{A} + \tilde{B}} = 1$. In particular, (11) fails to hold for \tilde{A} and \tilde{B} , but the good news is that Conjecture 1 does hold for them.

We note that $\tilde{B} = \tilde{A} + \tilde{A}$. Actually, if A is any set of three non-collinear points, and $B = A + A$, then there exists a linear transform φ such that A is a translate of $\varphi\tilde{A}$, and B is a translate of $\varphi\tilde{B}$. Therefore (11) does not hold for that A and B , as well. However, in the remainder of the paper, we prove the following theorem. From this result Conjecture 1 holds for the case when $|A| = b_A$ and $|B| = b_B$.

Theorem 6 *If A and B are finite non-collinear sets in \mathbb{R}^2 such that $|A| = b_A$, $|B| = b_B$ and (11) fails to hold, then either $|A| = 3$, and B is a translate of $A + A$, or $|B| = 3$, and A is a translate of $B + B$.*

To prove Theorem 6, we consider a unit vector v not parallel to any side of $[A]$ or $[B]$. We think of v as pointing vertically upwards. Let $l_{v,A}$ and $r_{v,A}$ be the leftmost and rightmost vertices of $[A]$, respectively. We note that $l_{v,A}$ and $r_{v,A}$ are unique, because v is not parallel to any side of $[A]$. Similarly, let $l_{v,B}$ and $r_{v,B}$ be the (unique) leftmost and rightmost vertices of $[B]$, respectively.

Remember that v points upwards. We observe that $l_{v,A}$ and $r_{v,A}$ divide the boundary of $[A]$ into an "upper" polygonal arc, and a "lower" polygonal arc. Let $A_{v,\text{upp}}$ and $A_{v,\text{low}}$ denote the set of points of A in the upper polygonal arc, and in the lower polygonal arc, respectively, excluding $l_{v,A}$ and $r_{v,A}$. For $a \in A$, we have

$$a \in A_{v,\text{upp}} \text{ iff } u \cdot v > 0 \text{ for any unit exterior normal } u \text{ to } [A] \text{ at } a; \quad (12)$$

$$a \in A_{v,\text{low}} \text{ iff } u \cdot v < 0 \text{ for any unit exterior normal } u \text{ to } [A] \text{ at } a. \quad (13)$$

In addition, as $l_{v,A}$ and $r_{v,A}$ are excluded, we have

$$|A_{v,\text{upp}}| + |A_{v,\text{low}}| = b_A - 2. \quad (14)$$

Similarly, $l_{v,B}$ and $r_{v,B}$ divide the boundary of $[B]$ into an "upper" polygonal arc, and a "lower" polygonal arc, and $B_{v,\text{upp}}$ and $B_{v,\text{low}}$ denote the set of points of B in the upper polygonal arc, and in the lower polygonal arc, respectively, excluding $l_{v,B}$ and $r_{v,B}$. For $b \in B$, we have

$$b \in B_{v,\text{upp}} \text{ iff } u \cdot v > 0 \text{ for any unit exterior normal } u \text{ to } [B] \text{ at } b; \quad (15)$$

$$b \in B_{v,\text{low}} \text{ iff } u \cdot v < 0 \text{ for any unit exterior normal } u \text{ to } [B] \text{ at } b; \quad (16)$$

$$|B_{v,\text{upp}}| + |B_{v,\text{low}}| = b_B - 2. \quad (17)$$

Lemma 7 *Let A and B be finite non-collinear sets in \mathbb{R}^2 , and let v be a unit vector not parallel to any side of $[A]$ or $[B]$. If $A_{v,\text{upp}}$, $A_{v,\text{low}}$, $B_{v,\text{upp}}$ and $B_{v,\text{low}}$ are all non-empty, then (11) holds.*

Proof: Lemma 2, (12) and (16) yield that $A_{v,\text{upp}} + B_{v,\text{low}} \subset \text{int}[A + B]$, therefore

$$i_{A+B} \geq |A_{v,\text{upp}} + B_{v,\text{low}}| \geq |A_{v,\text{upp}}| + |B_{v,\text{low}}| - 1.$$

In addition, Lemma 2, (13) and (15) yield that $A_{v,\text{low}} + B_{v,\text{upp}} \subset \text{int}[A + B]$, therefore

$$i_{A+B} \geq |A_{v,\text{low}} + B_{v,\text{upp}}| \geq |A_{v,\text{low}}| + |B_{v,\text{upp}}| - 1.$$

We deduce from (14) and (17) that

$$2i_{A+B} \geq |A_{v,\text{upp}}| + |B_{v,\text{low}}| + |A_{v,\text{low}}| + |B_{v,\text{upp}}| - 2 = b_A + b_B - 6.$$

□

In other words, Lemma 7 says that if (11) does not hold, then at least one of the sets $A_{v,\text{upp}}$, $A_{v,\text{low}}$, $B_{v,\text{upp}}$ and $B_{v,\text{low}}$ empty. We observe that replacing v by $-v$ simply exchanges $A_{v,\text{upp}}$ and $A_{v,\text{low}}$ on the one hand, and $B_{v,\text{upp}}$ and $B_{v,\text{low}}$ on the other hand. Therefore Proposition 9 will refine Lemma 7. Before that, we verify another auxiliary statement where $[p, q]$ denotes the closed line segment with end points $p, q \in \mathbb{R}^2$.

Lemma 8 *Let A and B be finite non-collinear sets in \mathbb{R}^2 , and let v be a unit vector not parallel to any side of $[A]$ or $[B]$. If $A_{v,\text{low}} = \emptyset$, then $i_{A+B} \geq |B_{v,\text{upp}}| - 2$, where equality yields that $B_{v,\text{low}} \subset [l_{v,B}, r_{v,B}]$, and the segments $[l_{v,A}, r_{v,A}]$ and $[l_{v,B}, r_{v,B}]$ are parallel.*

Proof: We drop the reference to v in the notation. After applying a linear transformation fixing v , we may assume that

$$w \cdot v = 0 \quad \text{for } w = r_A - l_A. \quad (18)$$

We may also assume that

$$l_A \cdot v = r_A \cdot v = 0. \quad (19)$$

If $r_B \cdot v > l_B \cdot v$, then we reflect both A and B through the line $\mathbb{R}v$. This keeps v , but interchanges the roles of l_A and r_A on the one hand, and the roles of l_B and r_B on the other hand. Therefore we may assume that

$$r_B \cdot v \leq l_B \cdot v. \quad (20)$$

Understanding exterior normals helps bound interior points in $[A + B]$. As A has some point above $[l_A, r_A]$ by $A_{\text{low}} = \emptyset$, (18) yields that

$$\text{either } u \cdot w > 0 \text{ or } u = -v \text{ for any exterior unit normal } u \text{ at } r_A \text{ to } [A], \quad (21)$$

$$\text{either } u \cdot w < 0 \text{ or } u = -v \text{ for any exterior unit normal } u \text{ at } l_A \text{ to } [A]. \quad (22)$$

We may assume that $B_{\text{upp}} \neq \emptyset$ (otherwise Lemma 8 trivially holds). We subdivide B_{upp} into the sets

$$B_{\text{upp}}^- = \{b \in B_{\text{upp}} : u \cdot w < 0 \text{ for any exterior unit normal } u \text{ at } b \text{ to } [B]\}, \quad (23)$$

$$B_{\text{upp}}^+ = \{b \in B_{\text{upp}} : u \cdot w > 0 \text{ for any exterior unit normal } u \text{ at } b \text{ to } [B]\}, \quad (24)$$

$$B_{\text{upp}}^0 = \{b \in B_{\text{upp}} : v \text{ is an exterior unit normal } u \text{ at } b \text{ to } [B]\}. \quad (25)$$

Since for any $b \in B$, the set of all exterior unit normals u at b to $[B]$ is an arc of the unit circle, the sets B_{upp}^- , B_{upp}^+ and B_{upp}^0 are pairwise disjoint, and their union is B_{upp} . In addition, we define

$$\tilde{B}_{\text{upp}}^- = \begin{cases} \{l_B\} \cup B_{\text{upp}}^- & \text{if there exists } b \in B \text{ with } b \cdot v < l_B \cdot v, \\ B_{\text{upp}}^- & \text{if } b \cdot v \geq l_B \cdot v \text{ for all } b \in B. \end{cases} \quad (26)$$

It follows that if $b \in \tilde{B}_{\text{upp}}^-$, then

$$\text{either } u \cdot w < 0 \text{ or } u = v \text{ for an exterior unit normal } u \text{ to } [B] \text{ at } b. \quad (27)$$

Turning to B_{upp}^0 , if $B_{\text{upp}}^0 \neq \emptyset$, then there exist $l_B^0, r_B^0 \in B_{\text{upp}}^0$ such that $r_B^0 - l_B^0 = sw$ for $s \geq 0$, and

$$B_{\text{upp}}^0 = B \cap [l_B^0, r_B^0], \quad (28)$$

$$v \cdot b_0 = \max\{v \cdot b : b \in B\} = H \text{ for all } b_0 \in B_{\text{upp}}^0. \quad (29)$$

To estimate i_{A+B} , we deduce from Lemma 2, and from (21) and (27) on the one hand, from (22) and (24) on the other hand, that

$$\begin{aligned} r_A + \tilde{B}_{\text{upp}}^- &\subset \text{int}[A + B] \text{ if } B_{\text{upp}}^- \neq \emptyset \\ l_A + B_{\text{upp}}^+ &\subset \text{int}[A + B] \text{ if } B_{\text{upp}}^+ \neq \emptyset. \end{aligned} \quad (30)$$

We claim that if $\tilde{B}_{\text{upp}}^- \neq \emptyset$ and $B_{\text{upp}}^+ \neq \emptyset$, then

$$\left| (r_A + \tilde{B}_{\text{upp}}^-) \cap (l_A + B_{\text{upp}}^+) \right| \leq 1. \quad (31)$$

We observe that $r_A + x = l_A + y$ if and only if $y - x = w$, and hence $x \cdot v = y \cdot v$. However, if $x_1, x_2 \in \tilde{B}_{\text{upp}}^-$ and $y_1, y_2 \in B_{\text{upp}}^+$ with $x_1 \cdot v = y_1 \cdot v < x_2 \cdot v = y_2 \cdot v$, then $(y_2 - x_2) \cdot w < (y_1 - x_1) \cdot w$, which in turn yields (31). We conclude by (19), (29), (30) and (31) then

$$|\{z \in (A + B) \cap \text{int}[A + B] : z \cdot v < H\}| \geq |\tilde{B}_{\text{upp}}^-| + |B_{\text{upp}}^+| - 1. \quad (32)$$

We recall that there exists some $p \in A_{\text{upp}}$, and hence $p \cdot v > 0$ by $l_A \cdot v = 0$. Thus if $B_{\text{upp}}^0 \neq \emptyset$, and $z \in \{l_A, r_A\} + B_{\text{upp}}^0$ is different from $l_A + l_B^0$ and $r_A + r_B^0$, then these two points of $A + B$ lie left and right from z . Since $(l_A + l_B) \cdot v < z \cdot v$, and $(p + l_B^0) \cdot v > z \cdot v$, we have $z \in \text{int}[A + B]$. In particular, $|\{l_A, r_A\} + B_{\text{upp}}^0| \geq |B_{\text{upp}}^0| + 1$ yields that

$$|\{z \in (A + B) \cap \text{int}[A + B] : z \cdot v = H\}| \geq |B_{\text{upp}}^0| - 1. \quad (33)$$

Adding (32) and (33) implies $i_{A+B} \geq |B_{\text{upp}}| - 2$. If $i_{A+B} = |B_{\text{upp}}| - 2$, then $\tilde{B}_{\text{upp}}^- = B_{\text{upp}}^-$, and hence $r_B \cdot v = l_B \cdot v$ by (20) and (26), and $B_{\text{low}} \subset [l_B, r_B]$. In particular (18) implies that $[l_{v,A}, r_{v,A}]$ and $[l_{v,B}, r_{v,B}]$ are parallel. \square

Proposition 9 *Let A and B be finite non-collinear sets in \mathbb{R}^2 , and let v be a unit vector not parallel to any side of $[A]$ or $[B]$. If (11) does not hold, then possibly after exchanging A and B , or v by $-v$, we have the following.*

- (i) $A_{v,\text{low}} = \emptyset$;
- (ii) $B_{v,\text{low}} \subset [l_{v,B}, r_{v,B}]$;
- (iii) $[l_{v,A}, r_{v,A}]$ and $[l_{v,B}, r_{v,B}]$ are parallel;
- (iv) either $B_{v,\text{low}} = \emptyset$ and $b_B = b_A$, or $|B_{v,\text{upp}}| = |A_{v,\text{upp}}| + |B_{v,\text{low}}| + 1$ and $b_B > b_A$.

Proof: We drop the reference to v in the notation. To present the argument, we make some preparations. Again using that (11) does not hold, Lemma 7 yields that possibly after exchanging A and B , or v by $-v$, we may assume that

$$A_{\text{low}} = \emptyset.$$

Possibly after exchanging A and B again, we may assume that

$$\text{if } B_{\text{low}} = \emptyset, \text{ then } b_B \geq b_A. \quad (34)$$

Since (11) does not hold, we have

$$i_{A+B} < \frac{1}{2}(b_A + b_B) - 3. \quad (35)$$

First we show that

$$\begin{aligned} & \text{either } |B_{\text{upp}}| = \frac{b_A + b_B}{2} - 2, \quad B_{\text{low}} = \emptyset \text{ and } b_A = b_B, \\ & \text{or } |B_{\text{upp}}| > \frac{b_A + b_B}{2} - 2. \end{aligned} \quad (36)$$

If $B_{\text{low}} = \emptyset$, then $b_B \geq b_A$ by (34), and hence

$$|B_{\text{upp}}| = b_B - 2 \geq \frac{b_A + b_B}{2} - 2,$$

with equality only if $b_A = b_B$.

If $B_{\text{low}} \neq \emptyset$, then we use that $A_{\text{upp}} \neq \emptyset$ by $A_{\text{low}} = \emptyset$. Thus Lemma 2, (12) and (16) yield that $A_{\text{upp}} + B_{\text{low}}$ lies in the interior of $[A + B]$. Combining this fact with (17) leads to

$$\begin{aligned} i_{A+B} & \geq |A_{\text{upp}} + B_{\text{low}}| \geq |A_{\text{upp}}| + |B_{\text{low}}| - 1 \\ & = b_A - 2 + b_B - 2 - |B_{\text{upp}}| - 1 = b_A + b_B - |B_{\text{upp}}| - 5. \end{aligned} \quad (37)$$

Therefore $|B_{\text{upp}}| > \frac{b_A + b_B}{2} - 2$ by (35), proving (36).

It follows from (35) and (36) that $i_{A+B} < |B_{\text{upp}}| - 1$, thus Lemma 8 implies that $i_{A+B} = |B_{\text{upp}}| - 2$, and in turn (ii) and (iii) of Proposition 9 hold. To prove (iv), we deduce from (35) that

$$b_A + b_B - 6 > 2i_{A+B} = 2|B_{\text{upp}}| - 4.$$

Therefore (36) yields that either $B_{\text{low}} = \emptyset$ and $b_A = b_B$, or

$$b_A + b_B - 4 < 2|B_{\text{upp}}| < b_A + b_B - 2.$$

In particular, $2|B_{\text{upp}}| = b_A + b_B - 3$ in the second case, which is in turn equivalent with $|B_{\text{upp}}| = |A_{\text{upp}}| + |B_{\text{low}}| + 1$ by $|A_{\text{upp}}| = b_A - 2$ and (17). In addition, $|B_{\text{upp}}| = |A_{\text{upp}}| + |B_{\text{low}}| + 1$ implies that $b_B > b_A$. \square

We have now developed enough machinery to prove Theorem 6. We repeat it here:

Theorem 10 *If A and B are finite non-collinear sets in \mathbb{R}^2 such that $|A| = b_A$, $|B| = b_B$ and (11) fails to hold, then either $|A| = 3$, and B is a translate of $A + A$, or $|B| = 3$, and A is a translate of $B + B$.*

Proof: We follow Proposition 9, and choose A , B , and v as in that result. For each $x \in A$, we have that if x lies on a corner of $[A]$, there exist vectors $v_{x,l}$ and $v_{x,r}$ such that $x = l_{v_{x,l},A}$ and $x = r_{v_{x,r},A}$. Since $A_{v,\text{low}} = \emptyset$, in the first case $r_{v_{x,l},A} = r_{v,A}$ and in the second $l_{v_{x,r},A} = l_{v,A}$. Consider one such

$x \in A_{v,upp}$. By Proposition 9, it follows that A is a subset of the triangle T_A formed by $l_{v,A}$, $r_{v,A}$, and x . And by the same proposition, all the sides of $[B]$ must be parallel to sides in A , so B is a subset of some triangle $T_B = \phi T_A$, where ϕ is a composition of a transposition and scalar multiplication. Then the corners of $[B]$ are $l_{v,B}$, $r_{v,B}$, and some point $y \in B$. We define the open line segments:

$$s_1 = (l_{v,A}, r_{v,A}) \quad (38)$$

$$s_2 = (l_{v,A}, x) \quad (39)$$

$$s_3 = (x, r_{v,A}) \quad (40)$$

$$t_1 = (l_{v,B}, r_{v,B}) \quad (41)$$

$$t_2 = (l_{v,B}, y) \quad (42)$$

$$t_3 = (y, r_{v,B}). \quad (43)$$

Let $A_i = s_i \cap A$ and $B_i = t_i \cap B$ for $i \in \{1, 2, 3\}$. Note that $A_1 = \emptyset$, and s_i is parallel to t_i , yet $A_i = \emptyset$ or $B_i = \emptyset$.

Assume for contradiction that $|A| > 3$. By Proposition 9, $|B_{v,upp}| \geq 2$. Thus $B_i \neq \emptyset$ for one $i \in \{2, 3\}$. Assume without loss of generality that $B_3 \neq \emptyset$; then by Proposition 9, $A_3 = \emptyset$ and so $A_2 \neq \emptyset$. Thus, letting $p \in A_2$, since B_1 and B_3 share no nonzero exterior normals with p , and since A_2 and $r_{v,B}$ share no nonzero exterior normals, $B_1 + p, A_2 + r_{v,B}, B_3 + p \in (A + B)_{int}$. And since $T_B = \phi T_A$, these three sets are pairwise disjoint. So

$$i_{A+B} \geq |B_1 + p| + |A_2 + r_{v,B}| + |B_3 + p| = b_A + b_B - 6, \quad (44)$$

and thus (11) holds, contrary to our assumption. So $|A| = 3$.

By Proposition 9, we have that if $B_{v,low} = \emptyset$, then $b_A = b_B = 3$. So, $2i_{A+B} \geq b_A + b_B - 6 = 0$, and again (11) holds. Thus, we have that $|B_{v,low}| \geq 1$, and so

$$|B_{v,upp}| = |B_{v,low}| + 2. \quad (45)$$

That is,

$$|B_2| + |B_3| = |B_1| + 1 \quad (46)$$

By the same argument, we get

$$|B_1| + |B_2| = |B_3| + 1 \quad (47)$$

$$|B_1| + |B_3| = |B_2| + 1 \quad (48)$$

It follows that $|B_1| = |B_2| = |B_3| = 1$ and so $b_B = 6$.

Now, $i_{A+B} > 0$, and if $i_{A+B} \geq 2$ then (11) holds, contradicting our assumption. Assuming then that $i_{A+B} = 1$, we let $b_i \in B_i$ for $i \in \{1, 2, 3\}$.

Then we see that $x + b_1 = r_{v,A} + b_2 = l_{v,A} + b_3$. And since $T_B = \phi T_A$, B must just be a translated version of $A + A$. And, as was mentioned in the beginning of this section, Conjecture 1 holds for A and B . \square

References

- [1] G.A. Freiman: Foundations of a structural theory of set addition. Translations of Mathematical Monographs, Vol 37. American Mathematical Society, Providence, R. I., 1973.
- [2] G.A. Freiman: Structure theory of set addition. II. Results and problems. In: Paul Erdős and his mathematics, I, 243-260, Bolyai Soc. Math. Stud., 11, János Bolyai Math. Soc., Budapest, 2002.
- [3] A. Matolcsi, I. Ruzsa (personal communication)
- [4] I.Z. Ruzsa: Sumsets and structure. In: Combinatorial number theory and additive group theory, Adv. Courses Math. CRM Barcelona, Birkhäuser Verlag, Basel, 2009, 87–210.
- [5] T. Tao, V. Vu: Additive combinatorics, Cambridge University Press, 2006.

K.J. Böröczky
Alfréd Rényi Institute of Mathematics
Hungarian Academy of Sciences
1053 Budapest, Reáltanoda u. 13-15
Hungary

B. Hoffman
Lewis & Clark College
615 Palatine Hill Road
Portland, OR 97219
United States