The minimum area of convex lattice n-gons

Imre Bárány *
Rényi Institute of Mathematics,
Hungarian Academy of Sciences
H-1364 Budapest Pf. 127 Hungary
barany@renyi.hu
and
Department of Mathematics
University College London
Gower Street, London, WC1E 6BT, UK

Norihide Tokushige
College of Education
Ryukyu University
Nishihara, Okinawa, 903-0213 Japan
hide@edu.u-ryukyu.ac.jp

June 25, 2001

Abstract

Let $A(n)$ be the minimum area of convex lattice n-gons. We prove that $\lim A(n)/n^3$ exists. Our computations suggest that the value of the limit is very close to $0.0185067\ldots$.

*Partially supported by Hungarian National Science Foundation Grants T 032452 and t 029255

1 Introduction

What is the minimal area $A(n)$ a convex lattice polygon with n vertices can have? The first to answer this question was G.E. Andrews [An63]. He proved that $A(n) \geq cn^3$ with some universal constant c. V.I. Arnol'd arrived to the same question from another direction [Ar80], and proved the same estimate. Further proofs are due to W. Schmidt [Schm85], Bárány-Pach [BP92]. The best lower bound comes form Rabinowitz [Ra93] via an inequality of Rényi-Sulanke [RS63]

$$\frac{1}{8\pi^2} < \frac{A(n)}{n^3} \leq \frac{1}{54}(1 + o(1)).$$

The upper bound follows from Remark 2 below.

Our main result is:

Theorem 1 $\lim A(n)/n^3$ *exists.*

The value of the limit — as we are going to show — equals the minimum of finitely many explicit extremal problems. But the finitely many is about 10^{10}, too many to solve. Our computations show, however, that most likely

$$\lim \frac{A(n)}{n^3} = 0.0185067\ldots.$$

We will also see that the convex lattice n-gon P with area $A(n)$ has elongated shape: after applying a suitable lattice preserving affine transformation, P has lattice width $c_1 n$ in direction $(0, 1)$ and has width $c_2 n^2$ in direction $(1, 0)$ where c_1, c_2 positive constants. Almost all the paper is devoted to the proof of Theorem 1.

Remark 1 Actually, Andrews [An63] showed much more, namely the following (see also [Schm85], [KS84]). If $P \subset \mathbf{R}^d$ is a convex lattice polytope with n vertices and volume $V > 0$, then

$$cn^{\frac{d+1}{d-1}} \leq V,$$

where c is a constant depending only on dimension.

2 Reduction

Define $\mathcal{P}_n$ as the set of all convex lattice n-gons in $\mathbf{R}^2$, then

$$A(n) = \min\{\text{Area}\, P : P \in \mathcal{P}_n\}.$$

In the next two claims, whose proof is given at the end of this section, we reduce the search for $A(n)$. As $A(n)$ is increasing it is enough to work with even n.

Claim 1 *For even n, there exists a centrally symmetric $P \in \mathcal{P}_n$ with $A(n) =$ Area P.*

Fix a centrally symmetric $P \in \mathcal{P}_n$ with $A(n) = \text{Area}\, P$ ($n = 2k$ even). The edges are $z_1, z_2, \dots, z_k, -z_1, \dots, -z_k$ in this order. Clearly, each z_i is a primitive vector, i.e., its components are coprime. Write $\mathbf{P}$ for the set of all primitive vectors in $\mathbf{Z}^2$. Define

$$C = \text{conv}\{z_1, z_2, \dots, z_k, -z_1, \dots, -z_k\}.$$

Then P is the zonotope spanned by $\{z_1, \dots, z_k\}$, i.e., $P = \sum_{i=1}^{k}[0, z_i]$ As it is well-known and easy to check

$$\text{Area}\, P(C) = \sum_{1 \le i < j \le k} |\det(z_i, z_j)|.$$

Write $\mathcal{C}$ for the set of 0-symmetric convex bodies in $\mathbf{R}^2$. Then $C \in \mathcal{C}$ and

$$A(C) = \frac{1}{8} \sum_{u \in C \cap \mathbf{P}} \sum_{v \in C \cap \mathbf{P}} |\det(u, v)|$$

which is clearly equal to Area $P(C)$.

Claim 2 *If $z \in C \cap \mathbf{P}$ then $z = z_i$, or $-z_i$ for some i.*

This means that the search for $A(n)$, or for minimal $P \in \mathcal{P}_n$ is reduced to the following minimization problem.

$$\text{Min}(n) = \min\{A(C) : C \in \mathcal{C} \text{ with } |C \cap \mathbf{P}| = n\}.$$

Observe that the solution C to the problem $\mathrm{Min}(n)$ is invariant under lattice preserving linear transformation. Thus we may fix C in *standard position*. This means that the lattice width (for the definition see [KL88]) of C is $2b = 2b(C)$ and is taken in direction $(0,1)$. Let $[-a,a]$ be the intersection of C with the x axis. We may further assume that the tangent line to C at $(a,0)$ has slope ≥ 1. A simple computation, (using the fact that the width of C in direction $(1,0)$ and $(1,-1)$ is at least $2b$) shows that $2a \geq b$. We fix C in this standard position. We record the following inequalities:

$$2a \geq b, \quad 2ab \leq \operatorname{Area} C \leq 4ab.$$

Remark 2 From now on we may assume $b \geq 2$ since for $b = 1$, according to Claim 2, the minimal C is (with $n = 2k$)

$$\mathrm{conv}\{\pm(0,1), \pm(1,1), \pm(2,1), \ldots, \pm(k-2,1), \pm(1,0)\}$$

which gives $\lim \frac{A(C)}{n^3} = \frac{1}{48}$, the example found in [Ra93]. When $C \in \mathcal{C}$ is a circle with $|C \cap \mathbf{P}| = n$, its radius, and then $A(C)$ are estimated easily showing $\lim \frac{A(C)}{n^3} = \frac{1}{54}$. This is the estimate given in the introduction.

Proof of Claim 1. Let $Q \in \mathcal{P}_n$ with vertices $v_1, v_2, \ldots, v_{2k}$ $(n = 2k)$ in this order. The diagonal $[v_i, v_{i+k}]$ cuts Q into two parts. Reflecting the part with smaller (or equal) area to the point $(v_i + v_{i+k})/2$ produces a lattice polygon with area $\leq \operatorname{Area} Q$. So it is enough to show that, for some $i \in \{1, \ldots, k\}$, the reflected n-gon is convex. It is certainly convex if there are parallel tangent lines to Q at v_i and v_{i+k}.

If there are no such tangents then the lines of the edges incident to v_i intersect those of incident to v_{i+k} on the same side of the line $v_i v_{i+k}$, on the left side, say. Then the lines of the edges, incident to v_{i+k} intersect those of v_{i+k+1} on the left side of the line $v_{i+1} v_{i+k+1}$, again. Starting with $i = 1$ a contradiction is reached at $i = k + 1$. ∎

We prove Claim 2 in stronger form:

Claim 2′ *Assume that $x_1, \ldots, x_k \in \mathbf{R}^2$, and no two of them collinear. If $x \in \mathrm{conv}\{\pm x_1, \ldots, \pm x_k\}$ and $x \neq \pm x_i$ $(\forall i)$, then there is a j such that replacing x_j by x gives a zonotope with smaller area.*

Proof. Assume first that x is on the boundary of $\operatorname{conv}\{\pm x_1,\dots,\pm x_k\}$. Then $x=(1-u)x_s+ux_t$ for some $0<u<1$. We may also assume that $\sum_{i=1}^k|\det(x_i,x_s)|\ge\sum_{i=1}^k|\det(x_i,x_t)|$. Then

$$\begin{aligned}
\sum_{i=1}^k|\det(x_i,x)| &= \sum_{i=1}^k|\det(x_i,(1-u)x_s+ux_t)|\\
&= \sum_{i=1}^k|(1-u)\det(x_i,x_s)+u\det(x_i,x_t)|\\
&< (1-u)\sum_{i=1}^k|\det(x_i,x_s)|+u\sum_{i=1}^k|\det(x_i,x_t)|\\
&\le (1-u)\sum_{i=1}^k|\det(x_i,x_s)|+u\sum_{i=1}^k|\det(x_i,x_s)|\\
&= \sum_{i=1}^k|\det(x_i,x_s)|.
\end{aligned}$$

Thus, replacing x_s by x makes the area smaller.

If x is in the interior of $\operatorname{conv}\{\pm x_1,\dots,\pm x_k\}$, then λx is on the boundary of this set with a unique $\lambda>1$ (apart from the trivial case $x=0$). The previous argument shows that replacing x_s by λx makes the area smaller, and consequently, replacing x_s by x makes it smaller, too. ∎

In the next two sections we approximate $|C\cap\mathbf{P}|$ and $A(C)$ using that the density of $\mathbf{P}$ in $\mathbf{Z}^2$ is $6/\pi^2$ (cf. [HW79]). We need to measure approximation by a quantity invariant under lattice preserving linear transformations. This is going to be the lattice width $2b=2b(C)$.

3 Approximating $|C\cap\mathbf{P}|$

Lemma 1

$$\left||C\cap\mathbf{P}|-\frac{6}{\pi^2}\operatorname{Area}C\right|\ll\operatorname{Area}C\cdot\frac{\log b}{b}.$$

Here and in what follows we use Vinogradov's $\ll$ notation. Thus $f(n)\ll g(n)$ means that $f(n)\le Dg(n)$ with some universal constant D.

Proof. The proof is standard and uses the Möbius function $\mu(d)$ see [HW79]. Set

$$C^+=C\cap\{(x,y)\in\mathbf{R}^2: y>0\}.$$

Clearly, $|C \cap \mathbf{P}| = 2 + 2|C^+ \cap \mathbf{P}|$ and

$$|C^+ \cap \mathbf{P}| = \sum_{(u,v)\in C^+\cap \mathbf{Z}^2} \sum_{\substack{d|u \\ d|v}} \mu(d) = \sum_{d=1}^{\infty} \mu(d) \sum_{\substack{(u,v)\in C^+\cap \mathbf{Z}^2 \\ d|u,\, d|v}} 1 =$$

$$= \sum_{d=1}^{b} \mu(d) |\frac{1}{d}C^+ \cap \mathbf{Z}^2|.$$

Claim 3

$$\left| |\frac{1}{d}C^+ \cap \mathbf{Z}^2| - \frac{1}{d^2}\mathrm{Area}\, C^+ \right| \le \frac{10a}{d}.$$

Proof. It suffices to show this for $d = 1$. Let $Q(z)$ denote the unit square centered at $z \in \mathbf{Z}^2$. Call $z \in \mathbf{Z}^2$ *inside* if $Q(z) \subset C^+$, *boundary* if $z \in C^+$ but $Q(z) \not\subset C^+$, and *outside* if $z \notin C^+$ and $Q(z) \cap \mathrm{int} C^+ \neq \emptyset$. Clearly

$$|C^+ \cap \mathbf{Z}^2| = |\{z \in \mathbf{Z}^2 : \text{inside}\}| + |\{z \in \mathbf{Z}^2 : \text{boundary}\}| =$$
$$= \mathrm{Area}\, C^+ + \sum_{z \text{ boundary}} \mathrm{Area}\,(Q(z) \setminus C^+) - \sum_{z \text{ outside}} \mathrm{Area}\,(Q(z) \cap C^+),$$

and the number of boundary and outside $z \in \mathbf{Z}^2$ is at most the perimeter of the smallest aligned box containing C^+, which is $2b + 2(a + b) \le 10a$. ∎

With Claim 3 we have

$$\begin{aligned} &\left| \sum_{d=1}^{b} \mu(d) \left(|\frac{1}{d}C^+ \cap \mathbf{Z}^2| - \frac{1}{d^2}\mathrm{Area}\, C^+ \right) \right| \\ &\le \sum_{d=1}^{b} \left| |\frac{1}{d}C^+ \cap \mathbf{Z}^2| - \frac{1}{d^2}\mathrm{Area}\, C^+ \right| \\ &\le \sum_{d=1}^{b} \frac{1}{d}(2b + 2(a+b)) \le 10a(1 + \log b). \end{aligned}$$

Thus

$$\left| |C^+ \cap \mathbf{P}| - \sum_{d=1}^{b} \frac{\mu(d)}{d^2}\mathrm{Area}\, C^+ \right| \le 10a(1 + \log b).$$

Note that $|\sum_{d=1}^{b} \frac{\mu(d)}{d^2} - \frac{6}{\pi^2}| \le \sum_{d=b+1}^{\infty} \frac{1}{d^2} < \frac{1}{b}$. Then

$$\begin{aligned}\left| |C \cap \mathbf{P}| - \frac{6}{\pi^2} \mathrm{Area}\, C \right| &\le 2 + \frac{1}{b} \mathrm{Area}\, C + 20a(1 + \log b) \\ &\le \mathrm{Area}\, C \left(\frac{2}{2ab} + \frac{1}{b} + \frac{10(1+\log b)}{b} \right) \\ &\ll \mathrm{Area}\, C \cdot \frac{\log b}{b}.\end{aligned}$$

This completes the proof of Lemma 1. ∎

4 Approximating $A(C)$

Lemma 2

$$\left| \sum_{u \in C^+ \cap \mathbf{P}} \sum_{v \in C^+ \cap \mathbf{P}} |\det(u,v)| - \left(\frac{6}{\pi^2}\right)^2 \int_{C^+} \int_{C^+} |\det(x,y)|\, dx\, dy \right| \ll (\mathrm{Area}\, C^+)^3 \frac{\log b}{b}.$$

Proof. It is very simple to see that

$$\det(u,v) = \int_{x \in Q(u)} \int_{y \in Q(v)} \det(x,y)\, dx\, dy.$$

Then this holds for $|\det(u,v)|$ with $|\det(x,y)|$ as the integrand if $\det(x,y)$ has constant sign on $Q(u) \times Q(v)$. This happens if $Q(u)$ and $Q(v)$ are separated by a line going through the origin. In case they are not separated, there are $\xi, \eta \in \mathbf{R}^2$ with $\|\xi\|_{\max}, \|\eta\|_{\max} \le \frac{1}{2}$ such that

$$0 = \det(u+\xi, v+\eta) = (u_1+\xi_1)(v_2+\eta_2) - (u_2+\xi_2)(v_1+\eta_1).$$

This shows that

$$\det(u,v) = u_1 v_2 - u_2 v_1 = -u_1\eta_2 - \xi_1 v_2 - \xi_1\eta_2 + u_2\eta_1 + v_1\xi_2 + \xi_2\eta_1$$

implying $|\det(u,v)| \le \frac{1}{2}(|u_1| + |u_2| + |v_1| + |v_2| + 1) \le \frac{1}{2}(2a + 2b + 1) \le 4a$ if $u, v \in \mathbf{Z}^2 - \{0\}$.

Similarly, if $Q(u)$ and $Q(v)$ are not separated, then for all $(x,y) \in Q(u) \times Q(v)$ (with $u, v \in \mathbf{Z}^2 - \{0\}$ again)

$$|\det(x,y)| \le 8a.$$

We start estimating $\sum\sum|\det(u,v)|$ via

$$\begin{aligned}
\sum_{u\in C^+\cap\mathbf{P}}\sum_{v\in C^+\cap\mathbf{P}}|\det(u,v)| &= \sum_{u\in C^+\cap\mathbf{Z}^2}\sum_{v\in C^+\cap\mathbf{Z}^2}|\det(u,v)|\sum_{s|u_1,s|u_2}\mu(s)\sum_{t|v_1,t|v_2}\mu(t)\\
&= \sum_{s=1}^{\infty}\sum_{t=1}^{\infty}\mu(s)\mu(t)\sum_{\substack{u\,\in\, C^+\cap\,\mathbf{Z}^2\\ s|u_1,\,s|u_2}}\sum_{\substack{v\,\in\, C^+\cap\,\mathbf{Z}^2\\ t|v_1,\,t|v_2}}|\det(u,v)|\\
&= \sum_{s=1}^{b}\sum_{t=1}^{b}s\mu(s)t\mu(t)\sum_{u\in\frac{1}{s}C^+\cap\mathbf{Z}^2}\sum_{v\in\frac{1}{t}C^+\cap\mathbf{Z}^2}|\det(u,v)|\\
&= \sum(1)-\sum(2)+\sum(3)
\end{aligned}$$

where

$$\sum(1)=\sum_s\sum_t s\mu(s)t\mu(t)\sum_{u\in\frac{1}{s}C^+\cap\mathbf{Z}^2}\sum_{v\in\frac{1}{t}C^+\cap\mathbf{Z}^2}\int_{Q(u)}\int_{Q(v)}|\det(x,y)|\,dx\,dy,$$

$\sum(2)$ is the same as $\sum(1)$ but for non-separated $Q(u),Q(v)$, and

$$\sum(3)=\sum_s\sum_t s\mu(s)t\mu(t)\sum_{u\in\frac{1}{s}C^+\cap\mathbf{Z}^2}\sum_{v\in\frac{1}{t}C^+\cap\mathbf{Z}^2}|\det(u,v)|\,dx\,dy,$$

again for non-separated $Q(u),Q(v)$. For fixed s and t, the number, $N(s,t)$, of non-separated pairs $Q(u),Q(v)$ with $u\in\frac{1}{s}C^+\cap\mathbf{Z}^2$ and $v\in\frac{1}{t}C^+\cap\mathbf{Z}^2$ can be estimated generously via Claim 3:

$$\begin{aligned}
N(s,t) &\le \left(\operatorname{Area}\frac{1}{s}C^+ + \frac{10a}{s}\right)\left(\operatorname{Area}\frac{1}{t}C^+ + \frac{10a}{t}\right)\\
&\ll (\operatorname{Area}C^+)^2\frac{1}{s^2t^2}.
\end{aligned}$$

In $\sum(3)$, $|\det(u,v)|\ll\frac{a}{s}+\frac{a}{t}$, and in $\sum(2)$ the integrand $|\det(x,y)|\ll\frac{a}{s}+\frac{a}{t}$ as well. Consequently

$$\begin{aligned}
|\sum(2)|,|\sum(3)| &\ll \sum_{s=1}^{b}\sum_{t=1}^{b}st\,(\operatorname{Area}C^+)^2\,\frac{1}{s^2t^2}\,\left(\frac{1}{s}+\frac{1}{t}\right)a\\
&\ll (\operatorname{Area}C^+)^3\,\frac{\log b}{b}.
\end{aligned}$$

Now we turn to $\sum(1)$. Define $R(s) = \bigcup_{u \text{ boundary}}(Q(u) \setminus \frac{1}{s}C^+)$ and $T(s) = \bigcup_{u \text{ outside}}(Q(u) \cap \frac{1}{s}C^+)$. We have to integrate over

$$\left[\frac{1}{s}C^+ \cup R(s) \setminus T(s)\right] \times \left[\frac{1}{t}C^+ \cup R(t) \setminus T(t)\right].$$

The main term comes from integrating over $\frac{1}{s}C^+ \times \frac{1}{t}C^+$. We are going to estimate the remaining 8 integrals.

It is readily seen that for $x, y \in C^+$ $|\det(x,y)| \le \operatorname{Area} C^+$. We need a slight strengthening of this (whose simple proof is omitted).

Claim 4 *When $x \in R(s) \cup T(s)$ and $y \in R(t) \cup T(t)$ and $s, t \le b$, then*

$$|\det(x,y)| \ll \frac{\operatorname{Area} C^+}{st}.$$

Using Claim 4

$$\int_{\frac{1}{s}C^+} \int_{R(t)} |\det(x,y)|\, dx\, dy \ll \frac{\operatorname{Area} C^+}{st} \int_{\frac{1}{s}C^+} dx \int_{R(t)} dy$$
$$\ll \frac{\operatorname{Area} C^+}{st} \frac{1}{s^2} \operatorname{Area} C^+ \frac{a}{t} \ll \frac{(\operatorname{Area} C^+)^3}{b} \frac{1}{s^3 t^2}.$$

So the sum of these terms multiplied by st is

$$\ll \sum_{s=1}^{b} \sum_{t=1}^{b} st \frac{(\operatorname{Area} C^+)^3}{b} \frac{1}{s^3 t^2} \ll (\operatorname{Area} C^+)^3 \frac{\log b}{b}.$$

The same applies to the integral over $\frac{1}{s}C^+ \times T(t)$ and when t and s are interchanged. Similarly

$$\int_{T(s)} \int_{T(t)} |\det(x,y)|\, dx\, dy \ll \frac{\operatorname{Area} C^+}{st} \frac{a}{s} \frac{a}{t} \ll \frac{(\operatorname{Area} C^+)^3}{b^2} \frac{1}{s^2} \frac{1}{t^2},$$

and the same works for the remaining three integrals. Thus we have

$$\left| \sum_{u \in C^+ \cap \mathbf{P}} \sum_{v \in C^+ \cap \mathbf{P}} |\det(u,v)| - \sum_{s=1}^{b} \sum_{t=1}^{b} s\mu(s) t\mu(t) \int_{\frac{1}{s}C^+} \int_{\frac{1}{t}C^+} |\det(x,y)|\, dx\, dy \right| \ll$$

$$\ll (\operatorname{Area} C^+)^3 \left(\frac{\log b}{b} + \frac{\log^2 b}{b^2}\right) \ll (\operatorname{Area} C^+)^3 \frac{\log b}{b}.$$

Here

$$
\begin{aligned}
&\sum_{s=1}^{b}\sum_{t=1}^{b} s\mu(s)t\mu(t)\int_{\frac{1}{s}C^+}\int_{\frac{1}{t}C^+} |\det(x,y)|\,dx\,dy \\
&\quad= \sum_{s=1}^{b}\sum_{t=1}^{b}\frac{\mu(s)}{s^2}\frac{\mu(t)}{t^2}\int_{C^+}\int_{C^+}|\det(x,y)|\,dx\,dy \\
&\quad= \left(\frac{6}{\pi^2}-\sum_{s=b+1}^{\infty}\frac{\mu(s)}{s^2}\right)\left(\frac{6}{\pi^2}-\sum_{t=b+1}^{\infty}\frac{\mu(t)}{t^2}\right)\int_{C^+}\int_{C^+}|\det(x,y)|\,dx\,dy.
\end{aligned}
$$

By Claim 4 $\int_{C^+}\int_{C^+}|\det(x,y)|\,dx\,dy \ll (\operatorname{Area} C^+)^3$. Thus

$$
\left|\sum_{s=1}^{b}\sum_{t=1}^{b} s\mu(s)t\mu(t)\int_{\frac{1}{s}C^+}\int_{\frac{1}{t}C^+}|\det(x,y)|\,dx\,dy - \left(\frac{6}{\pi^2}\right)^2\int_{C^+}\int_{C^+}|\det(x,y)|\,dx\,dy\right| \ll (\operatorname{Area} C^+)^3\frac{1}{b}
$$

finishing the proof of Lemma 2. ∎

5 Symmetrization

Theorem 2 *Assume $K \in \mathcal{C}$, and $B \in \mathcal{C}$ is a disk with* $\operatorname{Area} K = \operatorname{Area} B$. *Then*

$$
\int_K\int_K |\det(x,y)|\,dx\,dy \geq \int_B\int_B |\det(x,y)|\,dx\,dy.
$$

Equality holds iff K is an ellipsoid.

The proof goes by standard symmetrization (see e.g. [Schn93]). Let K^* be the symmetral of K with respect to a line l passing through O. We may assume, without loss of generality, that l is the x axis. To prove the theorem, it suffices to show the following:

Claim 5

$$
\int_K\int_K |\det(x,y)|\,dx\,dy \geq \int_{K^*}\int_{K^*} |\det(x,y)|\,dx\,dy.
$$

Proof. Fix $x_1 \in \mathbf{R}$ and set $K \cap \{(x_1, y) : y \in \mathbf{R}\} = [A, B]$. Then $K^* \cap \{(x_1, y) : y \in \mathbf{R}\} = [-(B-A)/2, (B-A)/2]$. Similarly fix y_1 and set $K \cap \{(x, y_1) : y \in \mathbf{R}\} = [C, D]$. The part of the integral $\int_K \int_K |\det(x, y)|\, dx\, dy$ with x_1, y_1 fixed and the same part of $\int_{K^*} \int_{K^*} |\det(x, y)|\, dx\, dy$ can be compared easily.

$$\int_C^D \int_A^B |x_1 y_2 - x_2 y_1|\, dx_2\, dy_2 \leq \int_{-(C-D)/2}^{(C-D)/2} \int_{-(A-B)/2}^{(A-B)/2} |x_1 y_2 - x_2 y_1|\, dx_2\, dy_2.$$

This is true since the integrand is the absolute value of a linear function on a rectangle $(x_2, y_2) \in [A, B] \times [C, D]$ which is clearly the smallest when the linear function is 0 at the center of the rectangle. This is the case exactly for the symmetral. ∎

If B is a disk centered at the origin, then

$$\int_B \int_B |\det(x, y)|\, dx\, dy = \frac{8}{9\pi^2} (\mathrm{Area}\, B)^3.$$

Thus Lemmas 1 and 2, and Theorem 2 give

Corollary 1 *If $C \in \mathcal{C}$ with $|C \cap \mathbf{P}| = n$, then*

$$A(C) \geq \left(\frac{1}{54} - D \frac{\log b}{b} \right) n^3$$

where D is a universal constant.

Remark 3 It turns out that one can take $D = 5000$ here when working with explicit constants instead of $\ll$.

6 The value of $\lim A(n)/n^3$

Set

$$f(z_1, \ldots, z_b) = \frac{2}{3} \sum_{i=1}^{b} \left(2\frac{\phi^2(i)}{i} + \frac{\phi(i)}{i^2} (\sum_{j=1}^{i-1} j\phi(j)) \right) z_i^3 + 2 \sum_{i=1}^{b} \phi(i) z_i \left(\sum_{j=1}^{i-1} \frac{\phi(j)}{j} z_j^2 \right).$$

We say that $z_1, \dots, z_b$ is *special* if the z_i are decreasing, $z_b \geq 0$ and they are convex, i.e., they satisfy $2z_i \geq z_{i-1} + z_{i+1}$ for $i = 2, \dots, b-1$. Define the following minimization problem:

$$\text{minimize } f(z_1, \dots, z_b) \text{ subject to } 4\sum_{i=1}^{b} \frac{\phi(i)}{i} z_i = 1, z_1, \dots, z_b \text{ is special.}$$

The minimum, which clearly exists, will be denoted by $M(b)$.

Theorem 3 $\lim A(n)/n^3 = \min_{b \leq 10^{10}} M(b)$.

Before the proof we give the following construction. Assume $b > 1$, and $x_1, \dots, x_b$ is special with $x_1 > 0$. Define

$$K = \operatorname{conv}\{(\pm x_i, \pm i) \in \mathbf{R}^2 : i = 1, \dots, b\}.$$

K is a convex set which is symmetric with respect to both axes. Define the 2 by 2 diagonal matrix H_n, with diagonal elements λ_n and 1, that satisfies $|\mathbf{P} \cap H_n K| = n$. (There might be a little ambiguity in this definition since several, but at most $4b$, elements may appear on the boundary of $H_n K$. Resolve it by considering some of these points as belonging, while some others as not belonging, to $H_n K$.) Setting $K_n = H_n K$ we see that $|\mathbf{P} \cap K_n| = n$.

Define $I(e, f; i) = \{(x, i) \in \mathbf{Z}^2 : e \leq x < f\}$. It is clear that the density of $\mathbf{P}$ on the line $y = i$ is $\phi(i)/i$, and there are exactly $\phi(i)$ primitive points on an interval of the form $I(si, (s+1)i; i)$. Write $I(i, s)$ for this interval. Now

$$|n - 4\sum_{i=1}^{b} \frac{\phi(i)}{i} \lambda_n x_i| \leq 4\sum_{1}^{b} i \leq 4b^2,$$

since, for each $i > 0$, the error term comes from the two subintervals $I_{left}(i)$ and $I_{right}(i)$ of $I(-\lambda_n x_i, \lambda_n x_i; i)$ that remain after deleting all $I(i, s)$ contained in it. A similar argument gives the following claim.

Claim 6

$$A(K_n) = \lambda_n^3 f(x_1, \dots, x_b) + O(\lambda_n^2),$$

where the implied constant depends only on b.

Proof. We only give a sketch. The basic observation is that $\sum |\det(u,v)|$ over $(u,v) \in (I(i,s) \cap \mathbf{P}) \times (I(j,t) \cap \mathbf{P})$ is the same as $\phi(i)\phi(j)/(ij)$ times the same sum over $(u,v) \in I(i,s) \times I(j,t)$, provided $\det(u,v)$ does not change sign on the box $I(i,s) \times I(j,t)$. The error terms come from two sources: First, from boxes where sign change occurs, but there are few of those, and there $|\det(u,v)| \le ij$. Secondly, when $u \in I_{left}(i)$ or $I_{right}(i)$, and similarly for v. But these intervals are short, with $|\det(u,v)|$ at most $4b\lambda_n$ (cf Claim 4). The statement follows by summing $\phi(i)\phi(j)|\det(u,v)|/(ij)$ over all $(u,v) \in (K_n \cap \mathbf{Z}^2) \times (K_n \cap \mathbf{Z}^2)$. ∎

So this construction satisfies $|\mathbf{P} \cap K_n| = n$ and

$$\lim \frac{A(K_n)}{n^3} = f(x_1, \ldots, x_b) \left(4 \sum_{i=1}^{b} \frac{\phi(i)}{i} x_i \right)^{-3}.$$

Proof of Theorem 1. Let C_n be the solution of the extremal problem $\mathrm{Min}(n)$ from Section 2. Let n_j be a sequence along which $\mathrm{Min}(n_j)$ tends to $M = \liminf \mathrm{Min}(n)$. If $b(C_n) \to \infty$ along a subsequence of n_j, then, according to Corollary 1

$$\lim \frac{A(C_n)}{n^3} = \frac{1}{54}$$

along the sequence n_j. But then this is true along the sequence n as well since, for the disk B_n containing n primitive points, $\lim A(B_n)/n^3 = 1/54$.

Assume now that $b(C_n)$ is bounded along n_j. Then we can choose a subsequence of n_j along which $b(C_n) = b$ for some fixed $b > 1$. To save writing, we denote this subsequence by n_j as well.

Let C_n^* denote the symmetral of C_n with respect to the y axis. The discrete analogue of the proof of Theorem 2 shows (we omit the straightforward details) that, along the sequence n_j,

$$\lim \frac{A(C_n)}{n^3} = \lim \frac{A(C_n^*)}{n^3}.$$

For $n = n_j$ and $i = 0, 1, \ldots, b$ define $x_i(n) \ge 0$ by

$$[-x_i(n), x_i(n)] = C_n^* \cap \{(x,i) : x \in \mathbf{R}\}.$$

Then with our previous notation $x_0(n) = a(C_n^*) = a(C_n)$, and $a(C_n) \to \infty$ (along n_j) since $\mathrm{Area}\, C_n \le 4a(C_n)b$. Choose now a subsequence of n_j along which $x_i(n)/a(C_n)$ is convergent, with limit x_i, for $i = 1, \ldots, b$. As the sequence $x_1, \ldots, x_b$ is special and $x_1 > 0$, the above construction works and gives the sequence K_n. It is obvious that, along the last subsequence of n_j, $\lim A(K_n)/n^3 = \lim A(C_n^*)/n^3 = M$. Then

$$\lim_{n\to\infty} \frac{A(K_n)}{n^3} = M$$

as well. ∎

Proof of Theorem 3. Given a special $x_1, \ldots, x_b$ with $4\sum_1^b \frac{\phi(i)}{i} x_i = 1$, we constructed a sequence of bodies K_n with $|K_n \cap \mathbf{P}| = n$ and $\lim A(K_n)/n^3 = f(x_1, \ldots, x_b)$. So the value of the limit in Theorem 1 is less than $1/54$ if we find a single special sequence on which f is smaller than $1/54$. Here is such a sequence with $b = 15$:

$$\begin{array}{lll}
x_1 = 0.03352589244, & x_2 = 0.03335447314, & x_3 = 0.03300806459, \\
x_4 = 0.03251038169, & x_5 = 0.03186074614, & x_6 = 0.03104245531, \\
x_7 = 0.03004944126, & x_8 = 0.02886386937, & x_9 = 0.02745127878, \\
x_{10} = 0.02577867736, & x_{11} = 0.02380388582, & x_{12} = 0.02143223895, \\
x_{13} = 0.01851220227, & x_{14} = 0.01470243266, & x_{15} = 0.008861427136,
\end{array}$$

giving

$$f(x_1, \ldots, x_{15}) = 0.0185067386955\ldots$$

which is smaller than $1/54 = 0.0185185185...$ by about 10^{-5}.

To see the bound $b \leq 10^{10}$, we use Corollary 1: if $b > 10^{10}$, and $C \in \mathcal{C}$ with $|C \cap \mathbf{P}| = n$, then

$$\frac{A(C)}{n^3} \geq \frac{1}{54} - 5000\frac{\log 10^{10}}{10^{10}} > 0.018507.$$

∎

Remark. The 10^{10} bound can be improved to about 10^7 by proving a stability version of Theorem 2: informally stated, this would say that if the left hand side of the inequality in Theorem 2 is smaller than $1 + \varepsilon$ times its right hand side, then K can be sandwiched between two ellipsoids E and $(1 + c\sqrt{\varepsilon})E$.

7 Remarks on computation

It seems hard to solve the minimization problems explicitly. We used the following heuristics. Let $x_1, \ldots, x_b$ be a solution to the problem. What can we expect about $x_1, \ldots, x_b$? According to Theorem 2, it is reasonable to assume that $(x_1, 1), \ldots, (x_b, b)$ are almost on the boundary of an ellipsoid. So let E_t be an ellipsoid whose half-axes are of length 1 and t with $b \le t \le b+1$, and define

$$w(i,t) = \sqrt{1 - \left(\frac{i}{t}\right)^2} \quad (0 \le i \le t).$$

Set $W = 4\sum_{i=1}^{b} \frac{\phi(i)}{i} w(i,t)$ and $z_i = w(i,t)/W$ for $i = 1, \ldots, b$. Then these $z_1, \ldots, z_b$ form a good approximation for the solution of the minimization problem. In fact, this method gives, with $t = 15.56$ (then $b = 15$) the points $z_1, \ldots, z_{15}$, that already satisfy $f(z_1, \ldots, z_{15}) < 1/54$. The even better solution $x_1, \ldots, x_{15}$ giving $f(x_1, \ldots, x_{15}) = 0.0185067\ldots$ was found near the previous $z_1, \ldots, z_{15}$ by solving the set of equations that constitute the necessary conditions for the extremum.

Using this heuristics we have checked the value of f near the ellipsoid E_t for $b = 1, \ldots, 100$ carefully (Figure 1) and for $b = 101, \ldots, 1000$ roughly (Figure 2). The computation suggests that the true limit of $A(n)/n^3$ is very close to the above value $0.0185067\ldots$. If this is the case then the minimizer P_n is very close, but not equal to, the ellipsoid with equation $x^2/A^2 + y^2/B^2 = 1$ where $A = 0.003573n^2$ and $B = 1.656n$. But even if the minimum is different, the shape of the minimizer P_n is oblong: it is $c_1 n$ wide $c_2 n^2$ long in its lattice width direction.

Figure 1



Figure 2



References

[An63] G. E. Andrews: *A lower bound for the volumes of strictly convex bodies with many boundary points.* Trans. AMS, **106** (1963), 270–279.

[Ar80] V. I. Arnol'd: *Statistics of integral convex polytopes.* (in Russian) Funk. Anal. Pril., **14** (1980), 1–3. English translation: Funct. Anal. Appl., **14** (1980), 79–84.

[BP92] I. Bárány, J. Pach: *On the number of convex lattice polygons.* Comb. Prob. Comp., **1** (1992), 295–302.

[HW79] G. H. Hardy, E. M. Wright: *An introduction to the theory of numbers,* Clarendon Press, Oxford (1979).

[KL88] R. Kanann, L. Lovász. *Covering minima and lattice-point-free convex bodies,* Annals of Math., **128** (1988) 577–602.

[KS84] S. B. Konyagin, K. A. Sevastyanov: *Estimation of the number of vertices of a convex integral polyhedron in terms of its volume.* (in Russian) Funk. Anal. Pril., **18** (1984), 13–15. English translation: Funct. Anal. Appl., **18** (1984), 11–13.

[Ra93] S. Rabinowitz: $O(n^3)$ *bounds for the area of a convex lattice* n*-gon.* Geombinatorics, **2** (1993), 85–88.

[RS63] A. Rényi, R. Sulanke: *Über die konvexe Hülle von* n *zufällig gewählten Punkten.* Z. Wahrsch. Verw. Geb., **2** (1963), 75-84.

[Schm85] W. Schmidt: *Integral points on surfaces and curves.* Monatshefte Math., **99** (1985), 45–82.

[Schn93] R. Schneider: *Convex bodies: the Brunn-Minkowski theory.* Encyclopedia Math. Appl. vol 44. Cambridge University Press, Cambridge, 1993.