

RESOLVABILITY AND MONOTONE NORMALITY

ISTVÁN JUHÁSZ, LAJOS SOUKUP, AND ZOLTÁN SZENTMIKLÓSSY

ABSTRACT. A space X is said to be κ -resolvable (resp. almost κ -resolvable) if it contains κ dense sets that are pairwise disjoint (resp. almost disjoint over the ideal of nowhere dense subsets). X is maximally resolvable iff it is $\Delta(X)$ -resolvable, where $\Delta(X) = \min\{|G| : G \neq \emptyset \text{ open}\}$.

We show that every crowded monotonically normal (in short: MN) space is ω -resolvable and almost μ -resolvable, where $\mu = \min\{2^\omega, \omega_2\}$. On the other hand, if κ is a measurable cardinal then there is a MN space X with $\Delta(X) = \kappa$ such that no subspace of X is ω_1 -resolvable.

Any MN space of cardinality $< \aleph_\omega$ is maximally resolvable. But from a supercompact cardinal we obtain the consistency of the existence of a MN space X with $|X| = \Delta(X) = \aleph_\omega$ such that no subspace of X is ω_2 -resolvable.

1. ω -RESOLVABILITY

For a topological space X we denote by $\mathcal{D}(X)$ the family of all dense subsets of X and by $\mathcal{N}(X)$ the ideal of all nowhere dense sets in X . Given a cardinal $\kappa > 1$, the space X is called κ -resolvable iff it contains κ many disjoint dense subsets. We say that X is *almost* κ -resolvable if there are κ many dense sets whose pairwise intersections are nowhere dense, that is we have $\{D_\alpha : \alpha < \kappa\} \subset \mathcal{D}(X)$ such that $D_\alpha \cap D_\beta \in \mathcal{N}(X)$ if $\alpha \neq \beta$. X is maximally resolvable iff it is $\Delta(X)$ -resolvable, where $\Delta(X) = \min\{|G| : G \neq \emptyset \text{ open}\}$ is called the dispersion character of X . Finally, if X is *not* κ -resolvable then it is also called κ -irresolvable.

The following simple but useful fact, in the case of κ -resolvability, was observed by El'kin in [4].

Lemma 1.1. *A space X is κ -resolvable (almost κ -resolvable) iff every nonempty open set in X includes a nonempty (and open) κ -resolvable (almost κ -resolvable) subset.*

2000 *Mathematics Subject Classification.* 54A35, 03E35, 54A25.

Key words and phrases. resolvable spaces, monotonically normal spaces.

The preparation of this paper was supported by OTKA grant no. 61600.

The aim of this paper is to present several results concerning the (almost) resolvability properties of monotonically normal spaces. Let us therefore recall their definition. For any topological space X we write

$$\mathcal{M}(X) = \{\langle x, U \rangle \in X \times \tau(X) : x \in U\}.$$

The elements of $\mathcal{M}(X)$ will be referred to as *marked* open sets. The space X is called *monotonically normal* iff it is T_1 and it admits a *monotone normality operator*, that is a function $H : \mathcal{M}(X) \longrightarrow \tau(X)$ such that

- (1) $x \in H(x, U) \subset U$ for each $\langle x, U \rangle \in \mathcal{M}(X)$,
- (2) if $H(x, U) \cap H(y, V) \neq \emptyset$ then $x \in V$ or $y \in U$.

We call a set D in a space X *strongly discrete* if the points in D may be separated by pairwise disjoint neighborhoods. It is well-known that in a monotonically normal space any discrete subset is strongly discrete. On the other hand, in [2] it was proved that every non-isolated point of a monotonically normal space is the accumulation point of a discrete subspace. Consequently, one obtains the following result.

Theorem 1.2 ([2]). *In a monotonically normal space any non-isolated point is the accumulation point of some strongly discrete set.*

Let us say that a space X is SD if it has the property described in theorem 1.2, that is every non-isolated point of X is the accumulation point of some strongly discrete set.

Theorem 1.3. *Any crowded SD space X is ω -resolvable.*

Proof. The SD property is clearly hereditary for open subspaces. Hence, by lemma 1.1, it suffices to prove that X includes an ω -resolvable subspace.

First we show that for every strongly discrete $D \subset X$ there is a strongly discrete $E \subset X \setminus \overline{D}$ such that $D \subset \overline{E}$. Indeed, fix a neighbourhood assignment U_d on D that separates D and for each $d \in D$ pick a strongly discrete set $E_d \subset X \setminus \{d\}$ with $d \in \overline{E_d}$. Then $E = \bigcup_{d \in D} (E_d \cap U_d)$ is clearly as claimed.

Now pick an arbitrary point $x \in X$ and set $E_0 = \{x\}$. Using the above claim, for each $n < \omega$ we can inductively define a strongly discrete set $E_{n+1} \subset X \setminus \overline{E_n}$ such that $E_n \subset \overline{E_{n+1}}$. Since $\bigcup_{i \leq n} E_i \subset \overline{E_n}$, the sets $\{E_n : n < \omega\}$ are pairwise disjoint. Let us finally set $E = \bigcup \{E_n : n < \omega\}$. It is clear from our construction that if $I \subset \omega$ is infinite then $\bigcup \{E_n : n \in I\}$ is dense in E , so the subspace E of X is obviously ω -resolvable. \square

Corollary 1.4. *Every crowded monotonically normal space is ω -resolvable.*

2. H-SEQUENCES AND ALMOST RESOLVABILITY

The main result of the previous section, namely that (crowded) monotonically normal spaces are ω -resolvable, used very little of the particular structure provided by monotone normality. In this section we shall describe a procedure on monotonically normal spaces that is quite specific in this respect and so, not surprisingly, it leads to some stronger (almost) resolvability results. This procedure had been originated (in a different form) by S. Williams and H. Zhou in [13].

Definition 2.1. Let H be a monotone normality operator on a space X . A family $\mathcal{E} \subset \mathcal{M}(X)$ of marked open sets is said to be *H-disjoint* if for any two members $\langle x, U \rangle, \langle y, V \rangle$ of \mathcal{E} we have $H(x, U) \cap H(y, V) = \emptyset$. Clearly, if \mathcal{E} is H-disjoint then $D(\mathcal{E}) = \{x : \exists U \text{ with } \langle x, U \rangle \in \mathcal{E}\}$ is (strongly) discrete.

By Zorn's lemma, for every open set G in X we can fix a *maximal* H-disjoint family $\mathcal{E}(G) \subset \mathcal{M}(G)$ with the additional property that $\overline{U} \subset G$ whenever $\langle x, U \rangle \in \mathcal{E}(G)$. The maximality of $\mathcal{E}(G)$ implies that

$$\bigcup \{H(x, U) : \langle x, U \rangle \in \mathcal{E}(G)\}$$

is a dense open subset of G .

With the help of the above defined operator $\mathcal{E}(G)$ we may now describe our basic procedure as follows.

Definition 2.2. A sequence $\langle \mathcal{E}_\alpha : \alpha < \delta \rangle$ is called a completed H-sequence of X iff

- (1) $\mathcal{E}_0 = \mathcal{E}(X)$,
- (2) for each $\alpha < \delta$ we have

$$\mathcal{E}_{\alpha+1} = \bigcup \{ \mathcal{E}(H(x, U) \setminus \{x\}) : \langle x, U \rangle \in \mathcal{E}_\alpha \},$$

- (3) if $\alpha < \delta$ is limit then the family

$$\mathcal{W}_\alpha = \{W \in \tau(X) : \forall \beta < \alpha \exists \langle x, U \rangle \in \mathcal{E}_\beta \text{ with } W \subset U\}$$

is a π -base in X (or, equivalently, its union $\bigcup \mathcal{W}_\alpha$ is dense in X) and \mathcal{E}_α is a maximal H-disjoint collection of marked open sets $\langle y, V \rangle$ with $V \in \mathcal{W}_\alpha$,

(4)

$$\mathcal{W}_\delta = \{W \in \tau(X) : \forall \beta < \delta \exists \langle x, U \rangle \in \mathcal{E}_\beta \text{ with } W \subset U\}$$

is *not* a π -base in X .

The reader may convince himself by a straight-forward transfinite induction that the following fact is valid.

Fact 2.3. *Every crowded monotonically normal space X , with monotone normality operator H , admits a completed H -sequence $\langle \mathcal{E}_\alpha : \alpha < \delta \rangle$ where δ is necessarily a limit ordinal.*

We now fix some notation concerning a given completed H -sequence $\langle \mathcal{E}_\alpha : \alpha < \delta \rangle$ of X . For any ordinal $\alpha < \delta$ we put $D_\alpha = D(\mathcal{E}_\alpha)$ and $H_\alpha = \bigcup \{H(x, U) : \langle x, U \rangle \in \mathcal{E}_\alpha\}$. It is clear from our definitions that each H_α is dense open in X , moreover $\beta < \alpha < \delta$ implies that $H_\beta \supset H_\alpha$ and $D_\beta \cap H_\alpha = \emptyset$. If $I \subset \delta$ is a set of ordinals we write $D[I] = \bigcup \{D_\alpha : \alpha \in I\}$. Finally, we set $V = X \setminus \overline{\bigcup \mathcal{W}_\delta}$, then V is a non-empty open set in X .

Lemma 2.4. *If I is bounded in δ then $D[I]$ is nowhere dense in X . However, if I is unbounded in δ then $D[I]$ is dense in V , that is $V \subset \overline{D[I]}$.*

Proof. The first part is obvious because $I \subset \alpha < \delta$ implies $D[I] \cap H_\alpha = \emptyset$.

Assume now that I is cofinal in δ but, arguing indirectly, for some $G \in \tau^*(V)$ we have $G \cap D[I] = \emptyset$. Pick any point $z \in G$, we claim that then, for all $\alpha < \delta$ and $\langle x, U \rangle \in \mathcal{E}_\alpha$, $H(x, U) \cap H(z, G) \neq \emptyset$ implies $z \in H(x, U)$.

Indeed, if $\beta \in (\alpha, \delta) \cap I$ then there is $\langle x', U' \rangle \in \mathcal{E}_\beta$ with

$$H(x', U') \cap H(x, U) \cap H(z, G) \neq \emptyset$$

because H_β is dense in X . It follows that $U' \subset H(x, U)$, hence $x' \notin G$ as $x' \in D_\beta$ and $G \cap D_\beta = \emptyset$. But then $H(x', U') \cap H(z, G) \neq \emptyset$ implies $z \in U' \subset H(x, U)$.

The sets $\{H(x, U) : \langle x, U \rangle \in \mathcal{E}_\alpha\}$ being pairwise disjoint, it follows that for each $\alpha < \delta$ there is exactly one $\langle x_\alpha, U_\alpha \rangle \in \mathcal{E}_\alpha$ such that $H(x_\alpha, U_\alpha) \cap H(z, G) \neq \emptyset$. But then $H(z, G) \subset \overline{H(x_\alpha, U_\alpha)} \subset \overline{U_\alpha}$ whenever $\alpha < \delta$, consequently

$$H(z, G) \subset \overline{U_{\alpha+1}} \subset U_\alpha$$

as well. This, however, would imply $H(z, G) \in \mathcal{W}_\delta$, contradicting that $H(z, G) \subset G \subset V$. \square

We may now give the main result of this section.

Theorem 2.5. *Any crowded monotonically normal space X is almost $\min(\mathfrak{c}, \omega_2)$ -resolvable. So X is almost ω_1 -resolvable, and even almost ω_2 -resolvable if the continuum hypothesis (CH) fails.*

Proof. By lemma 1.1 it suffices to show that some non-empty open $V \subset X$ satisfies this property. To see this, let us consider a completed H-sequence $\langle \mathcal{E}_\alpha : \alpha < \delta \rangle$ of X . Let I be a cofinal subset of δ of order type $\text{cf}(\delta)$ and $\{I_\zeta : \zeta < \mu\}$ be an almost disjoint subfamily of $[I]^{\text{cf}(\delta)}$, where $\mu = 2^\omega = \mathfrak{c}$ if $\text{cf}(\delta) = \omega$ and $\mu = \text{cf}(\delta)^+ \geq \omega_2$ if $\text{cf}(\delta) > \omega$. Then the family $\{D[I_\zeta] : \zeta < \mu\}$ witnesses that V is almost μ -resolvable. \square

Since almost ω -resolvability is clearly equivalent ω -resolvability, theorem 2.5 provides us a new proof of 1.4.

3. SPACES FROM TREES AND ULTRAFILTERS

The prime examples of monotonically normal spaces are metric and ordered spaces that are all known to be maximally resolvable. Compared to this the results of the two preceding sections seem rather modest. The main aim of this section is to show that, at least modulo some large cardinals, nothing stronger than ω -resolvability can be expected of a monotonically normal space X , even if the dispersion character $\Delta(X)$ is large. The examples that show this have actually been around but, as far as we know, the fact that they are monotonically normal has not been noticed.

The underlying set of such a space is an *everywhere infinitely branching* tree $\langle T, < \rangle$. This simply means that for each $t \in T$ the set S_t of all immediate successors of t in T is infinite. The height of such a tree is obviously a limit ordinal. (In fact, nothing is lost if we only consider trees of height ω .) By a *filtration* on T we mean a map F with domain T that assigns to every $t \in T$ a filter $F(t)$ on S_t such that every cofinite subset of S_t belongs to $F(t)$ (that is, $F(t)$ extends the Fréchet filter on S_t).

Definition 3.1. Assume that F is a filtration on an everywhere infinitely branching tree $\langle T, < \rangle$. A topology τ_F is then defined on T by

$$\tau_F = \{V \subset T : \forall t \in V \ (V \cap S_t \in F(t))\},$$

and the space $\langle T, \tau_F \rangle$ is denoted by $X(F)$.

Theorem 3.2. *Let F be a filtration on an everywhere infinitely branching tree $\langle T, < \rangle$. Then the space $X(F)$ is monotonically normal.*

Proof. That τ_F is indeed a topology that satisfies the T_1 separation axiom is obvious and well-known. The novelty is in showing that $X(F)$ is monotonically normal.

To this end we define $H(s, V)$ for $s \in V \in \tau_F$ as follows:

$$H(s, V) = \{t \in V : s \leq t \text{ and } [s, t] \subset V\}.$$

Of course, here $[s, t] = \{r : s \leq r \leq t\}$. Clearly, $H(s, V) \in \tau_F$ and $s \in H(s, V) \subset V$.

Next, assume that $t \in H(s_1, V_1) \cap H(s_2, V_2)$. Then $s_1, s_2 \leq t$ implies that s_1 and s_2 are comparable, say $s_1 \leq s_2$. But then we have $s_2 \in [s_1, t] \subset V_1$, consequently H is indeed a monotone normality operator on $X(F)$. \square

Of special interest are those filtrations F for which $F(t)$ is a (free) ultrafilter on S_t for all $t \in T$. Such an F will be called an *ultrafiltration*. In this case we have a convenient way to determine the closures of sets in the space $X(F)$ that will be put to good use later.

Definition 3.3. For every set $A \subset T$ we define

$$C(A) = A \cup \{t \in T : S_t \cap A \in F(t)\}.$$

Then by transfinite recursion we define $C^\alpha(A)$ for all ordinals α by $C^{\alpha+1}(A) = C(C^\alpha(A))$ for successors and $C^\alpha(A) = \bigcup\{C^\beta : \beta < \alpha\}$ for α limit.

Lemma 3.4. *Let F be an ultrafiltration on the tree T . Then a set $B \subset T$ is closed in $X(F)$ iff $B = C(B)$. Consequently, for any subset $A \subset T$ there is an ordinal $\alpha < |T|^+$ with $\overline{A} = C^\alpha(A)$.*

Proof. First, if $B = C(B)$ then for each $t \in T \setminus B$ we have $S_t \cap B \notin F(t)$, hence $S_t \setminus B \in F(t)$ because $F(t)$ is an ultrafilter. Then $T \setminus B$ is open by the definition of τ_F , hence B is closed. Conversely, if B is closed in $X(F)$ then for each $t \in T \setminus B$ we have $S_t \setminus B \in F(t)$, hence $S_t \cap B \notin F(t)$, that is $t \notin C(B)$. But this means that $B = C(B)$.

Next, $C(A) \subset \overline{A}$ is obvious, and then by induction we get $C^\alpha(A) \subset \overline{A}$ for all α . But for some $\alpha < |T|^+$ we must have $C(C^\alpha(A)) = C^\alpha(A)$, and then $\overline{A} = C^\alpha(A)$ for $C^\alpha(A)$ is closed by the above. \square

Let \mathfrak{u} be an ultrafilter on a set I and λ be a cardinal. \mathfrak{u} is said to be λ -*descendingly complete* iff $\bigcap\{X_\xi : \xi < \lambda\} \in \mathfrak{u}$ for each decreasing sequence $\{X_\xi : \xi < \lambda\} \subset \mathfrak{u}$. The ultrafilter \mathfrak{u} is called λ -*descendingly incomplete* iff it is not λ -descendingly complete. For example, \mathfrak{u} is countably complete exactly if it is ω -descendingly complete.

We shall need the following old result of Kunen and Prikry in our next irresolvability theorem for spaces obtained from certain ultrafiltrations.

Theorem (Kunen, Prikry, [9]). *If λ is a regular cardinal and \mathfrak{u} is a λ -descendingly complete ultrafilter (on any set) then \mathfrak{u} is also λ^+ -descendingly complete.*

Theorem 3.5. *Assume that F is an ultrafiltration on T and λ is a regular cardinal such that $F(t)$ is λ -descendingly complete for all $t \in T$. Then the space $X(F)$ is hereditarily λ^+ -irresolvable.*

Proof. First we show that for every set $A \subset T$ we have $\overline{A} = C^\lambda(A)$. By lemma 3.4 it suffices to show that $C(C^\lambda(A)) = C^\lambda(A)$.

Assume, indirectly, that $t \in C(C^\lambda(A)) \setminus C^\lambda(A)$, then we must have $C^\lambda(A) \cap S_t \in F(t)$. But

$$C^\lambda(A) \cap S_t = \bigcup_{\alpha < \lambda} C^\alpha(A) \cap S_t$$

where the right-hand side is an increasing union, hence there is an $\alpha < \lambda$ with $C^\alpha(A) \cap S_t \in F(t)$ because $F(t)$ is λ -descendingly complete. This implies that $t \in C^{\alpha+1}(A) \subset C^\lambda(A)$, a contradiction.

Let us now consider an *indexed* family of sets $\mathcal{F} = \{F_i : i \in I\}$. We are going to use the following notation:

$$\text{ord}(x, \mathcal{F}) = |\{i \in I : x \in F_i\}|$$

and

$$\text{ord}(\mathcal{F}) = \sup\{\text{ord}(x, \mathcal{F}) : x \in \bigcup_{i \in I} F_i\}.$$

Instead of the statement of the theorem we shall prove the following much stronger claim.

Lemma 3.6. *If $\mathcal{D} = \{D_i : i \in I\}$ is any indexed family of subsets of T with $\text{ord}(\mathcal{D}) \leq \lambda$ then $\text{ord}(\{\overline{D_i} : i \in I\}) \leq \lambda$ as well.*

Proof. We shall prove, by induction on $\alpha \leq \lambda$, that $\text{ord}(\mathcal{D}^\alpha) \leq \lambda$ where

$$\mathcal{D}^\alpha = \{C^\alpha(D_i) : i \in I\}.$$

We first show that $\text{ord}(\mathcal{D}^1) \leq \lambda$, this will clearly take care of all the successor steps.

Assume, indirectly, that $\text{ord}(t, \mathcal{D}^1) \geq \lambda^+$ for some $t \in T$, then we may find a set $J \in [I]^{\lambda^+}$ such that $t \in C(D_j) \setminus D_j$, hence $D_j \cap S_t \in F(t)$, for each $j \in J$.

By the theorem of Kunen and Prikry the ultrafilter $F(t)$ is also λ^+ -descendingly complete. Consequently, using a standard argument, one can show that there is an $L \in [J]^{\lambda^+}$ such that

$$\bigcap \{D_j \cap S_t : j \in L\} \neq \emptyset.$$

But this clearly contradicts $\text{ord}(\mathcal{D}) \leq \lambda$.

Next assume that $\alpha \leq \lambda$ is a limit ordinal and the inductive hypothesis holds for all $\beta < \alpha$. But now for each index $i \in I$ we have $C^\alpha(D_i) = \bigcup_{\beta < \alpha} C^\beta(D_i)$, hence

$$\text{ord}(t, \mathcal{D}^\alpha) \leq \sum_{\beta < \alpha} \text{ord}(t, \mathcal{D}^\beta) \leq |\alpha| \cdot \lambda = \lambda$$

whenever $t \in T$, and so $\text{ord}(\mathcal{D}^\alpha) \leq \lambda$. \square

It follows immediately from lemma 3.6 that if $\{A_i : i \in \lambda^+\}$ are pairwise disjoint non-empty subsets of T then the closures $\overline{A_i}$ cannot all be the same and so no subspace of $X(F)$ can be λ^+ -resolvable. \square

Corollary 3.7. *If F is an ultrafiltration on T such that $F(t)$ is countably complete for each $t \in T$ then $X(F)$ is ω -resolvable but hereditarily ω_1 -irresolvable. In particular, if κ is a measurable cardinal then there is a monotonically normal space X with $|X| = \Delta(X) = \kappa$ that is hereditarily ω_1 -irresolvable.*

The question if ω -resolvable spaces are also maximally resolvable was raised a long time ago by Ceder and Pearson in [1], and has just recently been settled completely in [7] (negatively). Corollary 3.7 yields a monotonically normal counterexample to this problem, from a measurable cardinal. Another counterexample from a measurable cardinal was found by Eckertson in [3], however, that example is not monotonically normal. We present two arguments to show this. First, Eckertson's example contains a crowded irresolvable subspace, hence it cannot be monotonically normal by corollary 1.4.

The second argument is based on our following observation that may have some independent interest. First we need some notation. If $\kappa \leq \lambda$ are cardinals we let τ_κ^λ denote the $< \kappa$ box product topology on 2^λ (generated by the base $\{[f] : f \in Fn(\lambda, 2; \kappa)\}$, where $[f] = \{x \in 2^\lambda : f \subset x\}$), moreover we set $\mathbb{C}_{\lambda, \kappa} = \langle 2^\lambda, \tau_\kappa^\lambda \rangle$.

Theorem 3.8. *If $\kappa^{<\kappa} = \kappa < \lambda$ then no dense subspace of $\mathbb{C}_{\lambda, \kappa}$ is monotonically normal.*

Proof of 3.8. Let X be dense in $\mathbb{C}_{\lambda, \kappa}$ and θ be a large enough regular cardinal. Let \mathcal{M} be an elementary submodel of $\langle \mathcal{H}(\theta), \in, \prec \rangle$ (where $\mathcal{H}(\theta)$ is the family of sets hereditarily of size $< \theta$ and \prec is a well-ordering of $\mathcal{H}(\theta)$) such that $|\mathcal{M}| = \kappa$ and $[\mathcal{M}]^{<\kappa} \subset \mathcal{M}$, moreover $X, \kappa, \lambda \in \mathcal{M}$. Note that then $Fn([\mathcal{M} \cap \lambda]^{<\kappa}, 2; \kappa) \subset \mathcal{M}$ as well.

Assume that X is monotonically normal and let $H \in \mathcal{M}$ be a monotone normality operator on X . We can assume that $H(x, [s] \cap X)$ is the trace on X of a basic open set for each basic open set $[s]$.

Let $I = \mathcal{M} \cap \lambda$ and pick $\alpha \in \lambda \setminus I$. $\mathcal{F} = \{f \restriction I : f \in \mathcal{M} \cap X\}$ is clearly dense in the subspace 2^I of $\mathbb{C}_{\lambda, \kappa}$. Let $\mathcal{F}_i = \{f \restriction I : f \in X \cap \mathcal{M} \wedge f(\alpha) = i\}$ for $i \in 2$ then $\mathcal{F} = \mathcal{F}_0 \cup \mathcal{F}_1$ so there is $i \in 2$ and $s \in Fn(I, 2; \kappa)$ such that \mathcal{F}_i is dense in $2^I \cap [s] \cap X$.

Let $b = s \cup \{\langle \alpha, 1 - i \rangle\}$ and pick $x \in X \cap [b]$. Next, let $H(x, [b] \cap X) = [b'] \cap X$ and $b'' = b' \restriction I$. Fix $b''' \in Fn(I, 2; \kappa)$ such that $b''' \supset b''$ and $x \notin [b''']$. Since \mathcal{F}_i is dense in $2^I \cap [s] \cap X$ we can pick $y \in X \cap \mathcal{M} \cap [b''']$ such that $y(\alpha) = i$. Let $[u] \cap X = H(y, [b'''] \cap X)$. Then $\text{dom } u \subset I$ because $H, b''', y \in \mathcal{M}$.

Since $x \notin [b''']$ and $y \notin [b]$ it follows that $H(x, [b]) \cap H(y, [b''']) = [u] \cap [b'] \cap X = \emptyset$. However $\text{supp } u \subset I$ and $u \supset b''' \supset b'' = b' \restriction I$, so u and b' are compatible functions of size $< \kappa$, i.e. $[u] \cap [b']$ is a nonempty open set in $\langle 2^\lambda, \tau_\kappa^\lambda \rangle$. Since X is dense we have $[u] \cap [b'] \cap X \neq \emptyset$, a contradiction. \square

Now, Eckertson's example obtained from a measurable cardinal κ contains a subspace homeomorphic to a dense subspace of $\mathbb{C}_{2^\kappa, \kappa}$, hence it cannot be monotonically normal by theorem 3.8 because $\kappa^{<\kappa} = \kappa$.

Of course, we have a space like in corollary 3.7 iff there is a measurable cardinal. Also, the cardinality (and dispersion character) of such a space is at least as large as the first measurable. But can one have a monotonically normal example that is much smaller? The answer to this question is, consistently, yes assuming the existence of a large cardinal that is even stronger than a measurable.

Theorem (Magidor, [11]). *It is consistent from a supercompact cardinal that there is an ω_1 -descendingly complete uniform ultrafilter on \aleph_ω .*

In [10] a similar statement was proved for $\aleph_{\omega+1}$ instead of \aleph_ω , but according to Magidor a slight modification of that proof works even for \aleph_ω .

From this result of Magidor and from theorem 3.5 we immediately obtain our promised result.

Corollary 3.9. *From a supercompact cardinal it is consistent to have a monotonically normal space X with $|X| = \Delta(X) = \aleph_\omega$ that is hereditarily ω_2 -irresolvable (hence not maximally resolvable).*

Actually, in [10] a slightly weaker result is given in which \aleph_ω is replaced with $\aleph_{\omega+1}$. However, in a private communication, Magidor pointed out to us that the method of [10] yields the above stronger version as well.

But can we do even better and go below \aleph_ω ? The answer to this question is, maybe surprisingly, negative. We are going to show that

any monotonically normal space of cardinality less than \aleph_ω is maximally resolvable. The proof of this result will be based on showing that all spaces of the form $X(F)$ with F an ultrafiltration on the tree $\text{Seq } \kappa = \kappa^{<\omega}$ of all finite sequences of ordinals less than κ are maximally resolvable provided that $\kappa < \aleph_\omega$. The first result to this effect, for constant ultrafiltrations on $\text{Seq } \omega_n$, was obtained by László Hegedüs in his Master's Thesis [5]. Of course, by a constant ultrafiltration we mean one for which $F(t)$ is the "same" ultrafilter for all $t \in T$.

Now, let κ be an arbitrary infinite cardinal. A non-empty subset T of $\text{Seq } \kappa$ is called a *subtree* of $\text{Seq } \kappa$ iff $t \restriction n \in T$ whenever $t \in T$ and $n < |t|$. For any subset A of $\text{Seq } \kappa$ we shall write $\min A$ to denote the set of all minimal elements of A (with respect to the tree ordering on $\text{Seq } \kappa$, of course).

If F is a filtration on $\text{Seq } \kappa$ and $v \in \text{Seq } \kappa$ we shall denote by F_v the derived filtration on $\text{Seq } \kappa$ defined by the formula $F_v(s) = F(v \frown s)$.

Assume now that S and $\{T_v : v \in \text{Seq } \kappa\}$ are subtrees of $\text{Seq } \kappa$. We then define their "sum" by

$$S \oplus \{T_v : v \in \text{Seq } \kappa\} = S \cup \{v \frown t : v \in \min(\text{Seq } \kappa \setminus S) \wedge t \in T_v\}.$$

Obviously, this sum is again a subtree of $\text{Seq } \kappa$.

If moreover f and $g = \{g_v : v \in \text{Seq } \kappa\}$ are functions with $\text{dom } f = S$ and $\text{dom } g_v = T_v$ then we define $f \oplus \{g_v : v \in \text{Seq } \kappa\} = f \oplus g$ by putting

$$\text{dom}(f \oplus g) = S \oplus \{T_v : v \in \text{Seq } \kappa\}$$

and

$$(f \oplus g)(x) = \begin{cases} f(x) & \text{for } x \in S \\ g_v(t) & \text{for } x = v \frown t \text{ with } v \in \min(\text{Seq } \kappa \setminus S), t \in T_v. \end{cases}$$

A subtree of $\text{Seq } \kappa$ is called *well-founded* iff it does not possess any infinite branches. Note that if S and $\{T_v : v \in \text{Seq } \kappa\}$ are all well-founded then so is $S \oplus \{T_v : v \in \text{Seq } \kappa\}$.

Now let $0 < \lambda \leq \kappa$ be cardinals and F be a filtration on $\text{Seq } \kappa$. We say that a function f is λ -good for F iff $\text{dom } f$ is a well-founded subtree of $\text{Seq } \kappa$, moreover $f[V] = \lambda$ whenever V is open in $X(F)$ with $\emptyset \in V$. As an easy (but useful) illustration of this concept we present the following result.

Lemma 3.10. *For each $0 < n < \omega$ and for any filtration F on κ there is a function f which is n -good for F .*

Proof. Let $\text{dom } f = \{s \in \text{Seq } \kappa : |s| < n\}$ and $f(s) = |s|$. □

The next result shows the relevance of these concepts to resolvability.

Theorem 3.11. *Let F be an filtration on $\text{Seq } \kappa$. If there are λ -good functions f_s for F_s for all $s \in \text{Seq } \kappa$ then $X(F)$ is λ -resolvable.*

Proof. Define the sequence of functions g_0, g_1, \dots by recursion as follows: $g_0 = f_\emptyset$ and $g_{n+1} = g_n \oplus \{f_s : s \in \text{Seq } \kappa\}$ for $n < \omega$. It is easy to check that then $g_\omega = \bigcup_{n < \omega} g_n$ maps $\text{Seq } \kappa$ to λ , i. e. $\text{dom } g_\omega = \text{Seq } \kappa$. Indeed, if $s \in \text{Seq } \kappa$ with $|s| = n$ then there is a $k \leq n$ with $s \in \text{dom } g_k$.

We show next that $g_\omega[V] = \lambda$ holds for any non-empty open set V in $X(F)$. Let n be such that $V \cap \text{dom } g_n \neq \emptyset$ and pick $v \in V \cap \text{dom } g_n$. Clearly, there is an extension s of v with $s \in V \cap \min(\text{Seq } \kappa \setminus \text{dom } g_n)$. Now let

$$W = \{t \in \text{Seq } \kappa : s \frown t \in V\}$$

then $\emptyset \in W$ and W is open in $X(F_s)$, hence $f_s[W] = \lambda$ because f_s is λ -good for F_s . But we clearly have $g_\omega(s \frown t) = f_s(t)$ for all $t \in \text{dom } f_s$, hence we have $g_\omega[V] = \lambda$ as well.

But then $\{g_\omega^{-1}(\alpha) : \alpha < \lambda\}$ is a pairwise disjoint family of dense sets in $X(F)$. \square

The following stepping-up type result will turn out to be very useful.

Lemma 3.12. *Assume that F is a filtration on $\text{Seq } \kappa$ such that $F(\emptyset)$ is λ -descendingly incomplete, moreover for every cardinal $\mu < \lambda$ and every ordinal $\alpha < \kappa$ there is a μ -good function f_μ^α for $F_{\langle \alpha \rangle}$. Then there is a λ -good function f for F .*

Proof. Fix a continuously decreasing sequence $\{X_\xi : \xi < \lambda\} \subset F(\emptyset)$ with empty intersection. For any ordinal $\nu < \lambda$ let us put $I_\nu = X_\nu \setminus X_{\nu+1}$, then we clearly have $\kappa = \bigcup \{I_\nu : \nu < \lambda\}$. For each $0 < \nu < \lambda$ fix a map $h_\nu : |\nu| \xrightarrow{\text{onto}} \nu$.

We now define the desired map f with the following stipulations:

$$\text{dom } f = \{\emptyset\} \cup \bigcup_{\nu < \lambda} \{\langle \alpha \rangle \frown t : \alpha \in I_\nu \text{ and } t \in \text{dom } f_{|\nu|}^\alpha\},$$

and for $s \in \text{dom } f$

$$f(s) = \begin{cases} 0 & \text{if } s = \emptyset, \\ h_\nu(f_{|\nu|}^\alpha(t)) & \text{if } s = \langle \alpha \rangle \frown t \text{ with } \alpha \in I_\nu, t \in \text{dom } f_{|\nu|}^\alpha. \end{cases}$$

Clearly, f is well-defined and $\text{dom } f$ is well-founded. If V is open in $X(F)$ with $\emptyset \in V$ then we have $V \cap S_\emptyset \in F(\emptyset)$ and hence

$$\sup\{\nu : \exists \alpha \in I_\nu \text{ with } \langle \alpha \rangle \in V\} = \lambda.$$

But $\langle \alpha \rangle \in V$ and $\alpha \in I_\nu$ imply $f_{|\nu|}^\alpha[\{s : \langle \alpha \rangle \frown s \in V\}] = |\nu|$ and so $f[V] \supset \nu$, hence we have $f[V] = \lambda$. \square

Theorem 3.13. *Let F be a filtration on $\text{Seq } \kappa$ and λ be an infinite cardinal such that $F(t)$ is μ -descendingly incomplete whenever $t \in \text{Seq } \kappa$ and $\omega \leq \mu \leq \lambda$. Then there are λ -good functions for all the derived filtrations F_s and hence $X(F)$ is λ -resolvable.*

Proof. The proof goes by a straight-forward induction on λ , using lemma 3.12 and the fact that our assumption on F is automatically valid also for all the derived filtrations F_s . The starting case $\lambda = \omega$ also uses lemma 3.10. The last statement is immediate from theorem 3.11. \square

A uniform ultrafilter on κ is trivially κ -descendingly incomplete. So if $\kappa = \omega_n < \aleph_\omega$, then it follows by n repeated applications of the above mentioned result of Kunen and Prikry that any uniform ultrafilter on κ is μ -descendingly incomplete for all μ with $\omega \leq \mu \leq \kappa$. Thus we get from theorem 3.13 the following result.

Corollary 3.14. *Assume that $\kappa < \aleph_\omega$ and F is any uniform ultrafiltration on $\text{Seq } \kappa$ (i. e. $F(t)$ is uniform for all $t \in \text{Seq } \kappa$). Then the space $X(F)$ is κ -resolvable.*

We now recall a definition from [8], see also [12].

Definition 3.15. Let X be a space and μ be an infinite cardinal number. We say that $x \in X$ is a T_μ point of X if for every set $A \in [X]^{<\mu}$ there is some $B \in [X \setminus A]^{<\mu}$ such that $x \in \overline{B}$. We shall use $T_\mu(X)$ to denote the set of all T_μ points of X .

The following result is an easy consequence of lemma 1.3 from [8]. In the particular case when μ is a successor cardinal it follows from proposition 2.1 of [12].

Lemma 3.16. *If $|X| = \mu$ is a regular cardinal and $T_\mu(X)$ is dense in X then X is μ -resolvable.*

This result will enable us to transfer certain results from spaces of the form $X(F)$, where F is a uniform ultrafiltration on $\text{Seq } \kappa$ for some regular cardinal κ , to monotonically normal and even more general spaces.

Let us recall from section 1 that every monotonically normal space is SD. In fact, as monotone normality is a hereditary property, it is even hereditarily SD (in short: HSD). We shall need below a property that is strictly between SD and HSD, namely that all *dense* subspaces are SD, we shall denote this property by DSD. It can be shown that for instance the Čech-Stone remainder ω^* is DSD but not HSD.

Theorem 3.17. *Assume that $\kappa = \text{cf}(\kappa) \geq \lambda$. Then the following are equivalent.*

- (1) *If X is a DSD space with $|X| = \Delta(X) = \kappa$ then X is λ -resolvable.*
- (2) *If X is a MN space with $|X| = \Delta(X) = \kappa$ then X is λ -resolvable.*
- (3) *If F is any uniform ultrafiltration on $\text{Seq } \kappa$ then the space $X(F)$ is λ -resolvable.*

Proof. Of course, only (3) \Rightarrow (1) requires proof. So assume (3) and consider a DSD space X with $|X| = \Delta(X) = \kappa$. If $T_\kappa(X)$ is dense in X then, by lemma 3.16 X is even κ -resolvable and we are done.

Otherwise, in view of lemma 1.1, we may assume that actually $T_\kappa(X) = \emptyset$. In this case for every point $x \in X$ there is a set $A_x \in [X]^{<\kappa}$ such that $x \in A_x$ and for $D_x = X \setminus A_x$ no $B \in [D_x]^{<\kappa}$ has x in its closure. Note that by $\Delta(X) = \kappa$ each D_x is dense in X .

But X is DSD, hence for every x there is a strongly discrete set $S_x \subset D_x$ with $x \in \overline{S_x}$. (Note that $S \subset D_x$ is strongly discrete in D_x iff it is so in X for D_x is dense.)

Next, by recursion on $|t|$, we define points x_t and open sets U_t in X as follows. First pick any point $x_\emptyset \in X = U_\emptyset$. If $x_t \in U_t$ has been defined then fix a one-to-one enumeration of $S_{x_t} \cap U_t = \{x_{t \smallfrown \alpha} : \alpha < \kappa\}$ and choose $\{U_{t \smallfrown \alpha} : \alpha < \kappa\}$ to be pairwise disjoint open neighbourhoods of them, all contained in U_t . Clearly, then the map $h : \text{Seq } \kappa \rightarrow X$ that maps t to $h(t) = x_t$ is injective.

Next, for any $t \in \text{Seq } \kappa$ extend the trace of the neighbourhood filter of x_t on $S_{x_t} \cap U_t$ to an ultrafilter u_t and define $F(t) = h^{-1}[u_t]$, which is an ultrafilter on $S_t = \{t \smallfrown \alpha : \alpha < \kappa\}$. It follows from our assumptions that every $F(t)$ is uniform and therefore $X(F)$ is λ -resolvable. But the subspace topology on $h[\text{Seq } \kappa]$ in X is clearly coarser than the h -image of τ_F , hence it is also λ -resolvable. By lemma 1.1, this completes our proof. \square

Corollary 3.18. *Let X be any DSD space of cardinality $< \aleph_\omega$. Then X is maximally resolvable. In particular, all MN spaces of size $< \aleph_\omega$ are maximally resolvable.*

Proof. Clearly, every open set U in X includes another open set V such that $|V| = \Delta(V)$. But every open subspace of a DSD space is again DSD, so theorem 3.17 and corollary 3.14 imply that V is $|V|$ -resolvable. But $\Delta(X) \leq |V|$, hence each such V is $\Delta(X)$ -resolvable and so, in view of lemma 1.1, X is maximally resolvable. \square

We conclude by listing a few open problems that we find interesting.

- Problem 3.19.** (1) *Is there a ZFC that example of a monotonically normal space that is not maximally resolvable?*
- (2) *Is it consistent to have a monotonically normal space X of cardinality less than the first measurable such that $\Delta(X) > \omega$ but X is not ω_1 -resolvable?*
- (3) *Is every crowded monotonically normal space almost \mathfrak{c} -resolvable?*

REFERENCES

- [1] J. Ceder and T. Pearson, *On products of maximally resolvable spaces*, Pacific J. Math. 22 1967 31–45.
- [2] Dow, A.; Tkachenko, M. G.; Tkachuk, V. V., and Wilson, R. G. *Topologies generated by discrete subspaces*, Glas. Mat. Ser. III 37(57) (2002), no. 1, 187–210.
- [3] Eckertson, F. W., *Resolvable, not maximally resolvable spaces*, Topology Appl. 79 (1997), no. 1, 1–11.
- [4] El’kin, A. G., *Resolvable spaces which are not maximally resolvable*, Vestnik Moskov. Univ. Ser. I Mat. Meh. 24 1969 no. 4 66–70.
- [5] Hegedűs, L., *Szűrők és lyukak*, Master’s Thesis, in Hungarian.
- [6] E. Hewitt, *A problem of set theoretic topology*, Duke Math. J. 10 (1943) 309–333.
- [7] I. Juhász, L. Soukup, and Z. Szentmiklóssy, *\mathcal{D} -forced spaces: a new approach to resolvability*, Top. Appl., 153 (2006), pp. 1800–1824.
- [8] I. Juhász, L. Soukup, and Z. Szentmiklóssy, *Resolvability of spaces having small spread or extent*, Top. Appl., to appear
- [9] Kunen, K. and Prikry, K., *On descendingly incomplete ultrafilters*, J. Symbolic Logic 36 (1971), 650–652.
- [10] Magidor, M., *On the singular cardinals problem. I*. Israel J. Math. 28 (1977), no. 1-2, 1–31.
- [11] Magidor, M., private communication.
- [12] O. Pavlov, *On resolvability of topological spaces*, Topology and its Applications 126 (2002) 37–47.
- [13] Williams, S. W. and Zhou, H. X., *Strong versions of normality*, in : General topology and applications (Staten Island, NY, 1989), 379–389, Lecture Notes in Pure and Appl. Math., 134, Dekker, New York, 1991.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS
E-mail address: juhasz@renyi.hu

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS
E-mail address: soukup@renyi.hu

EÖTVÖS UNIVERSITY OF BUDAPEST
E-mail address: zoli@renyi.hu